

Approximation of quadratic irrationals and their Pierce expansions

Jaume Paradís

Applied Mathematics, Univ. Pompeu Fabra
Rambla 32, 08002 Barcelona, Spain
e-mail: paradis-jaume@empr.upf.es

Lluís Bibiloni

Univ. Autònoma de Barcelona,
08193-Bellaterra, Barcelona, Spain
e-mail: ipdm0@cc.uab.es

Pelegri Viader

Applied Mathematics, Univ. Pompeu Fabra
Rambla 32, 08002 Barcelona, Spain
e-mail: viader_pelegri@empr.upf.es

June 12, 1996.

Abstract

In this article two aims are pursued: on the one hand, to present a rapidly converging algorithm for the approximation of square roots; on the other hand and based on the previous algorithm, to find the Pierce expansions of a certain class of quadratic irrationals as an alternative way to the method presented in 1984 by J.O. Shallit; we extend the method to find also the Pierce expansions of quadratic irrationals of the form $2(p-1)(p-\sqrt{p^2-1})$ which are not covered in Shallit's work.

1 Introduction.

In the year 1937, E.B. Escott published his paper *Rapid method for extracting a square root*, [4], where he presented an algorithm to find rational approximations for the square root of any real number. Escott's algorithm is based upon the algebraic identity:

$$\sqrt{\frac{x_1 + 2}{x_1 - 2}} = \frac{x_1 + 1}{x_1 - 1} \cdot \frac{x_2 + 1}{x_2 - 1} \cdot \frac{x_3 + 1}{x_3 - 1} \cdots,$$

where the x_i are obtained through the following recurrence:

$$x_n = x_{n-1}(x_{n-1}^2 - 3).$$

It is obvious that in order to calculate \sqrt{N} , Escott's algorithm must use rational x_i and thus the actual computation is considerably retarded.

More recently, in 1993, Y. Lacroix [5] refers to Escott's algorithm in the context of the representation of real numbers by generalized Cantor products and their metrical study.

In section 2 of this paper, we present an algorithm similar to Escott's but improved in the sense that we only use positive integers in the recurrence leading to the computation of \sqrt{N} . Moreover, the approximating fractions obtained by our algorithm constitute best approximations (of the second kind).

In 1984, J.O. Shallit [14] published the recurrence relations followed by the coefficients in the Pierce series development of irrational quadratics of the form $(c - \sqrt{c^2 - 4})/2$. Shallit's method is based upon Pierce's algorithm, [11], applied to the polynomial $x^2 - cx + 1$.

In section 3, we use the infinite product expansion provided by our square root algorithm to find the Pierce expansions, corresponding to irrationals of the form $p - \sqrt{p^2 - 1}$, as an alternative way to the one used by Shallit in [14]. The same method can also be used in the case of irrationals of the form $2(p - 1)(p - \sqrt{p^2 - 1})$ as we show in section 4.

2 The expansion of a quadratic irrational as an infinite product.

It is well-known (see [8, 9]) that the convergents p_n/q_n of the *regular continued fraction* development of \sqrt{r} , with r a positive integer, verify alternatively Pell's equations

$$p_n^2 - rq_n^2 = \pm 1,$$

and we get the recurrence relationships:

$$(1) \quad p_n = 2p_1p_{n-1} - p_{n-2}, \quad q_n = 2p_1q_{n-1} - q_{n-2},$$

that allow us to find all the solutions of Pell's equation from the first one (p_1, q_1) ; (we take $p_0 = 1, q_0 = 0$).

Lemma 2.1 *Let (p_1, q_1) be a positive solution ($p_1 > 0, q_1 > 0$) of Pell's equation $x^2 - ry^2 = 1$ where r is a positive integer free of squares. The sequence $\{(\bar{p}_n, \bar{q}_n)\}$ obtained recurrently in the way:*

$$(2) \quad \begin{cases} \bar{p}_n &= \bar{p}_{n-1}(4\bar{p}_{n-1}^2 - 3), & \bar{p}_1 &= p_1 \\ \bar{q}_n &= \bar{q}_{n-1}(4\bar{p}_{n-1}^2 - 1), & \bar{q}_1 &= q_1 \end{cases}$$

is a subsequence of the sequence $\{(p_n, q_n)\}$ of all solutions of the given Pell's equation, with the peculiarity that each solution is an integer multiple of the preceding.

Proof. We shall proceed by induction on n . Let us suppose that $\bar{p}_{n-1}^2 = r\bar{q}_{n-1}^2 + 1$ is verified. We must ascertain that:

$$(3) \quad \bar{p}_n^2 = r\bar{q}_n^2 + 1.$$

We replace \bar{p}_n and \bar{q}_n using the recurrence (2):

$$(4) \quad \bar{p}_{n-1}^2 (4\bar{p}_{n-1}^2 - 3)^2 = r\bar{q}_{n-1}^2 (4\bar{p}_{n-1}^2 - 1)^2 + 1.$$

To simplify let us denote by α the expression

$$\alpha = 4\bar{p}_{n-1}^2 - 2.$$

Equality (4) becomes:

$$\bar{p}_{n-1}^2(\alpha - 1)^2 = r\bar{q}_{n-1}^2(\alpha + 1)^2 + 1,$$

which can be written as

$$\bar{p}_{n-1}^2(\alpha^2 - 2\alpha + 1) = r\bar{q}_{n-1}^2(\alpha^2 + 2\alpha + 1) + 1.$$

Grouping together the terms corresponding to $\alpha^2 + 1$, we obtain:

$$(5) \quad (\alpha^2 + 1)(\bar{p}_{n-1}^2 - r\bar{q}_{n-1}^2) - 2\alpha(\bar{p}_{n-1}^2 + r\bar{q}_{n-1}^2) = 1,$$

and by the induction hypothesis (3),

$$\bar{p}_{n-1}^2 - r\bar{q}_{n-1}^2 = 1,$$

and also

$$\bar{p}_{n-1}^2 + r\bar{q}_{n-1}^2 = 2\bar{p}_{n-1}^2 - 1.$$

Thus equality (5) becomes

$$(6) \quad \alpha^2 - 2\alpha(2\bar{p}_{n-1}^2 - 1) = 0,$$

and, as we have $\alpha = 2(2\bar{p}_{n-1}^2 - 1)$, we deduce that (6) is in point of fact an algebraic identity. \diamond

Theorem 2.2 \sqrt{r} expands in an infinite product of the form:

$$\sqrt{r} = \frac{p_1}{q_1} \prod_{n=1}^{\infty} \frac{\alpha_n^2 - 3}{\alpha_n^2 - 1} = \frac{p_1}{q_1} \prod_{n=1}^{\infty} \left(1 - \frac{2}{\alpha_n^2 - 1} \right),$$

where (p_1, q_1) is a positive solution of Pell's equation $x^2 - ry^2 = 1$; $\alpha_1 = 2p_1$ and $\alpha_n = \alpha_{n-1}(\alpha_{n-1}^2 - 3)$.

Proof. With the same notations as in theorem 2.1 we have, on the one hand,

$$(7) \quad \sqrt{r} = \lim_{n \rightarrow \infty} \frac{\bar{p}_n}{\bar{q}_n},$$

and on the other we have the recurrence:

$$(8) \quad \frac{\bar{p}_n}{\bar{q}_n} = \frac{\bar{p}_{n-1}}{\bar{q}_{n-1}} \cdot \frac{4\bar{p}_{n-1}^2 - 3}{4\bar{p}_{n-1}^2 - 1}.$$

Iterating we obtain the expansion:

$$(9) \quad \frac{\bar{p}_n}{\bar{q}_n} = \frac{p_1}{q_1} \cdot \frac{4\bar{p}_1^2 - 3}{4\bar{p}_1^2 - 1} \cdot \frac{4\bar{p}_2^2 - 3}{4\bar{p}_2^2 - 1} \cdots \frac{4\bar{p}_{n-1}^2 - 3}{4\bar{p}_{n-1}^2 - 1},$$

or, if we prefer it, we can simplify expression (9) defining the new recurrence:

$$(10) \quad \alpha_n = \alpha_{n-1}(\alpha_{n-1}^2 - 3), \quad \alpha_1 = 2p_1,$$

which allow us to write:

$$(11) \quad \frac{\bar{p}_n}{\bar{q}_n} = \frac{p_1}{q_1} \cdot \frac{\alpha_1^2 - 3}{\alpha_1^2 - 1} \cdots \frac{\alpha_{n-1}^2 - 3}{\alpha_{n-1}^2 - 1}.$$

Finally, taking limits as $n \rightarrow \infty$, the expansion of \sqrt{r} in an infinite product is:

$$(12) \quad \sqrt{r} = \frac{p_1}{q_1} \prod_{n=1}^{\infty} \frac{\alpha_n^2 - 3}{\alpha_n^2 - 1} = \frac{p_1}{q_1} \prod_{n=1}^{\infty} \left(1 - \frac{2}{\alpha_n^2 - 1} \right). \quad \diamond$$

Using the recurrence (10) in (11) we obtain:

$$(13) \quad \frac{\bar{p}_n}{\bar{q}_n} = \frac{\alpha_n}{2q_1(\alpha_1^2 - 1) \cdots (\alpha_{n-1}^2 - 1)}.$$

The recurrence (10) is a fast way to compute the fractions of (13) which constitute best approximations of the second kind of any irrational quadratic of the form \sqrt{r} where r is a positive integer; to start, we just need a positive solution of Pell's equation $x^2 - ry^2 = 1$. With ten iterations of the algorithm we obtain a fraction whose approximation to the irrational is of the order $10^{-30,000}$. With 14 iterations the approximation gives us a million correct decimal figures.

Expansion (12), among others, is the one considered by Y. Lacroix in [5], in connection with Cantor's representation of real numbers by infinite products (see [1]).

3 The Pierce expansion of $p - (p^2 - 1)^{1/2}$.

Any real number $\alpha \in (0, 1]$ has a unique Pierce expansion of the form:

$$(14) \quad \alpha = \frac{1}{a_1} - \frac{1}{a_1 a_2} + \cdots + \frac{(-1)^{n+1}}{a_1 a_2 \cdots a_n} + \cdots,$$

where $\{a_n\}$ is a strictly increasing sequence of positive integers. These a_i will be called *coefficients* or *partial quotients* of the development.

Following Erdős and Shallit [3] we will denote the right hand side of (14) by the special symbol:

$$\langle a_1, a_2, \dots, a_n, \dots \rangle.$$

If expansion (14) is infinite, then α is irrational. Otherwise α is rational.

One of the first mathematicians to consider these developments was Lambert in [7]. Later, Lagrange refers to them in [6], but we have to wait to Sierpinski, [16], and Ostrogadsky, [12], (independently) to see their numerical properties studied. Perron mentions them in [10] among other unusual series representations of real numbers. T. A. Pierce in [11] used them to approximate roots of algebraic equations, and quite recently, in 1986, J. O. Shallit [15] studied their metrical properties using methods developed by Rényi in [13] to study the metrical properties of Engel's series (series of the type (14) but with all its signs positive, see [2, 10, 13]). The same Shallit, using Pierce's algorithm obtained in 1984, see [14], the Pierce expansion of all irrational quadratics of the form

$$(15) \quad \frac{c - \sqrt{c^2 - 4}}{2} \quad \text{with integer } c, c \geq 3.$$

If $c = 2k$, the irrational in (15) is directly of the form $k - \sqrt{k^2 - 1}$. If $c = 2k + 1$ it can be seen that the irrational in (15) can be written as:

$$\frac{1}{2k} - \frac{1}{2k(2k+2)} + \frac{1}{2k(2k+2)} \cdot (p - \sqrt{p^2 - 1}) \quad \text{with } p = (2k+1)(2k^2 + 2k - 1).$$

Thus, Pierce expansion of irrationals of the form studied by Shallit are a particular case of irrationals of the form $p - \sqrt{p^2 - 1}$. The aim of this section is to find the Pierce expansion of all irrationals of the form $p - \sqrt{p^2 - 1}$.

Now, if $\sqrt{p^2 - 1} = q\sqrt{r}$ with r free of squares, (p, q) is a solution of Pell's equation $x^2 - ry^2 = 1$.

Theorem 3.1 *Given r , a positive integer free of squares, let (p, q) be a positive solution of Pell's equation $x^2 - ry^2 = 1$. The Pierce expansion of the irrational $p - q\sqrt{r}$ is exactly*

$$(16) \quad p - q\sqrt{r} = \langle \alpha_1 - 1, \alpha_1 + 1, \alpha_2 - 1, \alpha_2 + 1, \dots \rangle$$

where $\alpha_1 = 2p$, and $\alpha_{n+1} = \alpha_n(\alpha_n^2 - 3)$.

Proof. Using expression (13):

$$\frac{\bar{p}_n}{\bar{q}_n} = \frac{\alpha_n}{2q_1(\alpha_1^2 - 1) \cdots (\alpha_{n-1}^2 - 1)},$$

we can write its right hand side as:

$$\frac{\alpha_{n-1}}{2q_1(\alpha_1^2 - 1) \cdots (\alpha_{n-2}^2 - 1)} \cdot \frac{\alpha_{n-1}^2 - 3}{\alpha_{n-1}^2 - 1} = \frac{\alpha_{n-1}}{2q_1(\alpha_1^2 - 1) \cdots (\alpha_{n-2}^2 - 1)} \left(1 - \frac{2}{\alpha_{n-1}^2 - 1}\right).$$

Now, as we have the algebraic identity:

$$\frac{\alpha_{n-1}}{b(\alpha_{n-1}^2 - 1)} = \frac{1}{b(\alpha_{n-1} - 1)} - \frac{1}{b(\alpha_{n-1} - 1)(\alpha_{n-1} + 1)},$$

iterating the former process we eventually reach the expansion:

$$\begin{aligned} \frac{\bar{p}_n}{\bar{q}_n} &= \frac{p_1}{q_1} - \frac{1}{q_1(\alpha_1 - 1)} + \frac{1}{q_1(\alpha_1 - 1)(\alpha_1 + 1)} + \cdots + \\ &+ \frac{1}{q_1(\alpha_1^2 - 1) \cdots (\alpha_{n-2}^2 - 1)(\alpha_{n-1} - 1)} - \frac{1}{q_1(\alpha_1^2 - 1) \cdots (\alpha_{n-1}^2 - 1)}. \end{aligned}$$

In our case, $p_1 = p$ and $q_1 = q$ and we can write:

$$\begin{aligned} \frac{p}{q} - \frac{p_n}{q_n} &= \frac{1}{q(\alpha_1 - 1)} - \frac{1}{q(\alpha_1 - 1)(\alpha_1 + 1)} + \cdots + \\ (17) \quad &+ \frac{1}{q(\alpha_1^2 - 1) \cdots (\alpha_{n-2}^2 - 1)(\alpha_{n-1} - 1)} - \frac{1}{q(\alpha_1^2 - 1) \cdots (\alpha_{n-1}^2 - 1)}. \end{aligned}$$

As $n \rightarrow \infty$ we obtain the infinite Pierce expansion:

$$(18) \quad \frac{p}{q} - \sqrt{r} = \sum_{i=1}^{\infty} \left(\frac{1}{q \prod_{k=1}^{i-1} (\alpha_k^2 - 1) \cdot (\alpha_i - 1)} - \frac{1}{q \prod_{k=1}^i (\alpha_k^2 - 1)} \right),$$

which is equivalent to (16). \diamond

4 The Pierce expansion of $2(p-1)[p - (p^2 - 1)^{1/2}]$.

In this section we are going to see how the method we have just used can be extended to find the Pierce expansion of irrational quadratics of the form $2(p-1)(p - \sqrt{p^2 - 1})$.

As above, our starting point will be Pell's equation $x^2 - ry^2 = 1$, and we will choose a subsequence of the sequence of its solutions. We will need the following result:

Lemma 4.1 *Given a positive solution (p, q) of Pell's equation $x^2 - ry^2 = 1$, with r free of squares, the recurrent sequence $\{(\bar{p}_n, \bar{q}_n)\}$ obtained in the form:*

$$\begin{cases} \bar{p}_n &= 2\bar{p}_{n-1}^2 - 1, & \bar{p}_1 &= p \\ \bar{q}_n &= 2\bar{p}_{n-1}\bar{q}_{n-1}, & \bar{q}_1 &= q \end{cases}$$

is a subsequence of the sequence $\{(p_n, q_n)\}$ of all the equation solutions.

Proof. The result is easily proved by induction. Let us suppose that $\bar{p}_{n-1}, \bar{q}_{n-1}$ verify $\bar{p}_{n-1}^2 - r\bar{q}_{n-1}^2 = 1$. For the next index we will have:

$$\begin{aligned}\bar{p}_n^2 &= 2\bar{p}_{n-1}^2 - 1 = 4\bar{p}_{n-1}^4 - 4\bar{p}_{n-1}^2 + 1 \\ r\bar{q}_n^2 &= r4\bar{p}_{n-1}^2\bar{q}_{n-1}^2,\end{aligned}$$

and subtracting:

$$\bar{p}_n^2 - r\bar{q}_n^2 = 4\bar{p}_{n-1}^2(\bar{p}_{n-1}^2 - r\bar{q}_{n-1}^2) - 4\bar{p}_{n-1}^2 + 1 = 1. \quad \diamond$$

Once proved that all pairs (\bar{p}_n, \bar{q}_n) are solutions of the given Pell's equation and using that

$$\sqrt{r} = \lim_{n \rightarrow \infty} \frac{\bar{p}_n}{\bar{q}_n},$$

we will try, as before, to expand the fraction \bar{p}_n/\bar{q}_n as a finite Pierce expansion, and then, taking limits, obtain the infinite Pierce expansion corresponding to the irrational \sqrt{r} , or an equivalent one.

Let us start with the fraction \bar{p}_n/\bar{q}_n , and let us express its numerator and denominator in terms of the preceding pair of solutions:

$$(19) \quad \frac{\bar{p}_n}{\bar{q}_n} = \frac{2\bar{p}_{n-1}^2 - 1}{2\bar{p}_{n-1}\bar{q}_{n-1}} = \frac{\bar{p}_{n-1}}{\bar{q}_{n-1}} - \frac{1}{2\bar{p}_{n-1}\bar{q}_{n-1}}.$$

Proceeding with the expansion of the equation above, we will eventually reach the first one, \bar{p}_1/\bar{q}_1 and the chain of equalities:

$$\begin{aligned}\frac{\bar{p}_n}{\bar{q}_n} &= \frac{\bar{p}_1}{\bar{q}_1} - \frac{1}{2\bar{p}_1\bar{q}_1} - \frac{1}{2\bar{p}_2\bar{q}_2} - \dots - \frac{1}{2\bar{p}_{n-1}\bar{q}_{n-1}} = \\ &= \frac{\bar{p}_1}{\bar{q}_1} - \frac{1}{\bar{q}_1 2\bar{p}_1} - \frac{1}{\bar{q}_1 2\bar{p}_1 2\bar{p}_2} - \dots - \frac{1}{\bar{q}_1 2\bar{p}_1 2\bar{p}_2 \dots 2\bar{p}_{n-1}} = \\ &= \frac{\bar{p}_1}{\bar{q}_1} - \frac{1}{\bar{q}_1} \left(\frac{1}{2\bar{p}_1} + \frac{1}{2\bar{p}_1 2\bar{p}_2} + \dots + \frac{1}{2\bar{p}_1 2\bar{p}_2 \dots 2\bar{p}_{n-1}} \right).\end{aligned}$$

Taking limits in this last expression, and remembering that $\bar{p}_1 = p$ and $\bar{q}_1 = q$ we obtain:

$$(20) \quad \sqrt{r} = \frac{p}{q} - \frac{1}{q} \left(\frac{1}{2p} + \frac{1}{2p2\bar{p}_2} + \dots + \frac{1}{2p2\bar{p}_2 \dots 2\bar{p}_{n-1}} + \dots \right),$$

where the \bar{p}_i follow the recurrence

$$\bar{p}_n = 2\bar{p}_{n-1}^2 - 1, \quad \bar{p}_1 = p.$$

The series within the parenthesis in the right hand side of (20) is an Engel's series.

Equality (20) can also be expressed in the form:

$$(21) \quad p - q\sqrt{r} = \frac{1}{2p} + \frac{1}{2p\bar{p}_2} + \dots + \frac{1}{2p2\bar{p}_2 \dots 2\bar{p}_{n-1}} + \dots$$

or even, if we prefer it, we can state the result:

Lemma 4.2 For all positive integers, p , we have:

$$(22) \quad p - \sqrt{p^2 - 1} = \frac{1}{2\bar{p}_1} + \frac{1}{2\bar{p}_1\bar{p}_2} + \cdots + \frac{1}{2\bar{p}_1 2\bar{p}_2 \cdots 2\bar{p}_{n-1}} + \cdots$$

with $\bar{p}_i = 2\bar{p}_{i-1}^2 - 1$, $\bar{p}_1 = p$.

Expression (22) is known as Stratemeyer's formula, and can be obtained algebraically by the method described in Perron's [10, Ch. IV]. We mention in passing that the recurrence (22) verified by the \bar{p}_i is exactly the recurrence verified by the denominators in the infinite product expansion presented by Cantor in [1].

Now we are ready for the following result:

Theorem 4.3 If p is a positive integer greater than one,

$$2(p-1)(p - \sqrt{p^2 - 1}) = \langle 1, p_1, p_2, p_3, \cdots \rangle,$$

where $p_1 = p$ and the p_i verify:

$$\begin{cases} p_{2n} &= 4(p_{2n-1} + 1) \\ p_{2n+1} &= 2p_{2n-1}^2 - 1. \end{cases}$$

Proof. To prove theorem 4.3 we just have to change the Engel's series in (22) into a Pierce expansion. In order to do that let us consider the Pierce expansion of theorem 4.3:

$$(23) \quad \langle 1, p_1, p_2, p_3, \cdots \rangle$$

with the recurrence

$$\begin{aligned} p_{2n} &= 4(p_{2n-1} + 1) \\ p_{2n+1} &= 2p_{2n-1}^2 - 1, \quad p_1 = p. \end{aligned}$$

If we denote by S the irrational number represented by (23) we have the following expansion

$$\begin{aligned} S &= 1 - \frac{1}{p_1} + \frac{1}{p_1 p_2} - \cdots = \\ &= \frac{p_1 - 1}{p_1} + \frac{p_3 - 1}{p_1 p_2 p_3} + \cdots + \frac{p_{2n+1} - 1}{p_1 p_2 \cdots p_{2n+1}}. \end{aligned}$$

We want to see that each fraction in the sum above is of the form:

$$(24) \quad \frac{p_{2n+1} - 1}{p_1 p_2 \cdots p_{2n+1}} = \frac{p_1 - 1}{p_1 2 p_3 \cdots 2 p_{2n+1}}.$$

We will proceed by induction on n . For $n = 0$ it is trivially true. Let us expand the left hand side of (24) in the following way:

$$(25) \quad \frac{p_{2n+1} - 1}{p_1 \cdots p_{2n-1} p_{2n} p_{2n+1}} = \underbrace{\frac{p_{2n-1} - 1}{p_1 \cdots p_{2n-1}}}_{(*)} \cdot \underbrace{\left(\frac{p_{2n+1} - 1}{p_{2n+1} p_{2n}} \right)}_{(**)} \cdot \frac{1}{p_{2n-1} - 1}.$$

The term $(\star\star)$ can be written as follows:

$$\frac{2p_{2n-1}^2 - 1 - 1}{p_{2n+1}4(p_{2n-1} + 1)} \cdot \frac{1}{p_{2n-1} - 1} = \frac{2(p_{2n-1}^2 - 1)}{p_{2n+1}4(p_{2n-1}^2 - 1)} = \frac{1}{2p_{2n+1}}.$$

Now, by the induction hypothesis applied to factor (\star) in (25), we obtain finally

$$(26) \quad \frac{p_{2n+1} - 1}{p_1 \cdots p_{2n-1} p_{2n} p_{2n+1}} = \frac{p_1 - 1}{p_1 2p_3 \cdots 2p_{2n-1}} \cdot \frac{1}{2p_{2n+1}}.$$

Thus S can be written as:

$$\begin{aligned} S &= \langle 1, p_1, p_2, \cdots \rangle = \\ &= \frac{p_1 - 1}{p_1} + \frac{p_1 - 1}{p_1 2p_3} + \cdots + \frac{p_1 - 1}{p_1 2p_3 \cdots 2p_{2n+1}} + \cdots = \\ &= \frac{p_1 - 1}{p_1} \left(1 + \frac{1}{2p_3} + \frac{1}{2p_3 2p_5} + \cdots + \frac{1}{2p_3 2p_5 \cdots 2p_{2n+1}} + \cdots \right) = \\ &= 2(p_1 - 1) \underbrace{\left(\frac{1}{2p_1} + \frac{1}{2p_1 2p_3} + \cdots + \frac{1}{2p_1 2p_3 \cdots 2p_{2n+1}} \right)}_{(\star\star\star)}. \end{aligned}$$

But by Stratemeyer's formula (22) the term $(\star\star\star)$ is precisely $p_1 - \sqrt{p_1^2 - 1}$. This ends the proof of theorem 4.3. \diamond

5 Conclusions

The algorithm presented in this article provides fast best approximations to any irrational of the form \sqrt{r} , where r is a positive integer. At the same time, the algorithm provides the necessary background to obtain the Pierce expansion of some quadratic irrationals whose partial quotients, a_i , grow as x^3 . The procedure used proves also that the convergents in the Pierce expansions of these irrationals are best approximations of the second kind.

We also present the Pierce series development of the irrationals of the form $2(p-1)(p - \sqrt{p^2 - 1})$, whose partial quotients grow as x^2 .

There exist though quadratic irrationals that escape the above laws, and whose partial quotients obey the metrical behaviour $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = e$, found by J.O Shallit in [15].

References

- [1] G. Cantor. "Zwei Sätze über eine gewisse Zerlegung der Zahlen in unendliche Produkte". *Schlömilch, Zeitschrift f. Math. u. Physik.* **14**, pp. 152–158, 1869.
- [2] P. Erdős, A. Rényi, and P. Szűsz. "On Engel's and Sylvester's series". *Ann. Univ. Sci. Budapest. Sect. Math.* **1**, pp. 7–32, 1958.

- [3] P. Erdős, J.O. Shallit. “New bounds on the Length of Finite Pierce and Engel Series”. *Séminaire de Théorie de Nombres Bordeaux*. **3**, pp. 43–53, 1991.
- [4] E.B. Escott. “Rapid method for extracting a square root”. *Amer. Math. Monthly*. **44**, pp. 644–646, 1937.
- [5] Y. Lacroix. “Metric properties of generalized Cantor products”. *Acta Arithmetica*. LXIII.1, pp. 61–77, 1993.
- [6] J.L. Lagrange. “Essai d’analyse numérique sur la transformation des fractions”. *Journal de l’École Polytechnique*, Vème cahier, t. II, prairial, an VI. Also in *Oeuvres* ed. J.A. Serret), Tome VII, Section quatrième, pp. 291–313.
- [7] J.H. Lambert. *Verwandlung der Brüche, Beiträge zum Gebrauche der Mathematik und deren Anwendung*. 1770.
- [8] I. Niven, H.S. Zuckerman, H.L. Montgomery. *An Introduction to the Theory of Numbers*. Wiley & Sons Inc. N.Y. 5th ed. 1991.
- [9] C.D. Olds. *Continued Fractions*. The Mathematical Association of America, 1963.
- [10] O. Perron. *Irrationalzahlen*. De Gruyter, Berlin u. Leipzig, 2nd. ed, 1939.
- [11] T.A. Pierce. “On an algorithm and its use in approximating roots of algebraic equations”. *Amer. Math. Monthly*. **36**, pp. 523–525, 1929.
- [12] E. Ya. Remez. “On series with alternating signs, which may be related to two algorithms of M. V. Ostrogradski for the approximation of irrationals numbers”. *Uspehi Matem. Nauk. (N.S.)*, **6,5** (45), pp. 33–42, 1951.
- [13] A. Rényi. “A new approach to the theory of Engel’s series”. *Ann. Univ. Sci. Budapest Sec. Math.* **5** pp. 25–32, 1962.
- [14] J.O. Shallit. “Some predictable Pierce Expansions”. *Fibonacci Quart.* **22**, pp. 332–335, 1984.
- [15] J.O. Shallit. “Metric theory of Pierce Expansions”. *Fibonacci Quart.* **24**, pp. 22–40, 1986.
- [16] W. Sierpinski. “Sur quelques algorithmes pour développer les nombres réels en séries” (1911). Included in *Oeuvres choisies*. t.1 pp. 236–254, 1974.

1991 *Mathematics Subject Classification*. Primary 11A67, 65D99.

Keywords and phrases. Quadratic irrationals, Pierce series, Square root extraction.