

# ALMOST SURE TESTABILITY OF CLASSES OF DENSITIES

Luc Devroye  
School of Computer Science  
McGill University  
Montreal, Canada H3A 2A7

Gábor Lugosi  
Department of Economics  
Pompeu Fabra University  
Ramon Trias Fargas, 25-27  
08005 Barcelona, Spain

ABSTRACT. Let a class  $\mathcal{F}$  of densities be given. We draw an i.i.d. sample from a density  $f$  which may or may not be in  $\mathcal{F}$ . After every  $n$ , one must make a guess whether  $f \in \mathcal{F}$  or not. A class is almost surely testable if there exists such a testing sequence such that for any  $f$ , we make finitely many errors almost surely. In this paper, several results are given that allow one to decide whether a class is almost surely testable. For example, continuity and square integrability are not testable, but unimodality, log-concavity, and boundedness by a given constant are.

KEYWORDS AND PHRASES. Density estimation, kernel estimate, convergence, testing, asymptotic optimality, minimax rate, minimum distance estimation, total boundedness.

1991 MATHEMATICS SUBJECT CLASSIFICATIONS: Primary 62G05.

RUNNING HEAD: TESTING DENSITIES

---

The first author's work was supported by NSERC Grant A3456 and by FCAR Grant 90-ER-0291. The second authors' work was supported by DIGES Grant PB96-0300.

## Table of Contents

1. Introduction
2. Classes defined by functionals
3. Classes with vanishing minimax risk
4. Testability via kernel estimates
5. Testability via total boundedness
6. Testability via Yatracos' minimum distance estimates
7. Smoothness and monotonicity classes
8. Ad hoc analysis: testing unimodality and convexity
9. Proof of Theorem 2
10. Non-testable classes
11. Non-testability of mixture classes

## 1. Introduction

Let  $\mathcal{F}$  be a class of densities on the real line. Assume that the sequence of i.i.d. random variables  $X_1, X_2, \dots$  is drawn according to some density  $f$  which may, or may not belong to  $\mathcal{F}$ . The question is if it is possible to detect from a finite sample whether the underlying density is a member of the class  $\mathcal{F}$ . Examples of  $\mathcal{F}$  include the class of all normal densities, the class of all densities bounded by 5, the class of all unimodal densities, or the class of densities supported on  $[0, 5]$  which are Lipschitz with a Lipschitz constant less than 2. We investigate the testability of such properties.

A **test** is a sequence  $\{T_n\}$  of functions  $T_n : \mathcal{R}^n \rightarrow \{0, 1\}$  so that upon observing the sample  $X_1, \dots, X_n$ , one guesses that the unknown density  $f$  is in  $\mathcal{F}$  iff  $T_n(X_1, \dots, X_n) = 1$ . A class  $\mathcal{F}$  is called almost surely testable, or simply **testable**, if there exists a test such that for any density  $f$

$$\mathbf{P} \{T_n(X_1, \dots, X_n) \neq I_{\{f \in \mathcal{F}\}} \text{ for only finitely many } n\} = 1.$$

(Here  $I$  denotes the indicator function.) In other words, we require that the test make the right decision eventually, almost surely, for any density. A test  $\{T_n\}$  with the above property is called **consistent**. Obviously,  $\mathcal{F}$  is testable if and only if its complement  $\mathcal{F}^c$  is testable.

This paper points out that many common assumptions one finds nowadays in papers simply are not testable. For example, one cannot test that a density has compact support or is bounded. One cannot even test whether it is Lipschitz, let alone have a continuous first derivative. Integrability conditions like  $\int f^2 < \infty$  can also not be tested. On the other hand, many properties can be tested such as unimodality, monotonicity, and convexity. Several general theorems in this respect are provided. For example, closed classes with an  $L_1$  minimax risk over  $\mathcal{F}$  that tends to zero are testable. We provide an explicit and non-obvious construction of such tests.

Even though we work with densities on the real line, basically all results may be reproduced for multivariate densities in  $\mathcal{R}^d$ . Since the multivariate problem does not need new ideas, we stay with the simple case of  $d = 1$ .

Finally, we recall here that Kulkarni and Zeitouni (1995) studied a similar, though slightly weaker definition of testability. Their definition, based on early work of Cover (1973) and Koplowitz (1977), is asymmetric. If  $f \notin \mathcal{F}$  they allow the test to fail for some very small subclass of densities. Under their definition, Kulkarni and Zeitouni propose a general testing procedure. This, however, cannot be used to prove testability under the symmetric definition studied here.

## 2. Classes defined by functionals

In this section we derive some sufficient conditions for testability for classes which are defined in terms of some functional of the density. A functional  $\Psi$  assigns an (extended) real number to every density. Examples include  $\Psi(f) = \int f^2$ ;  $\Psi(f) = \int f \log f$ ;  $\Psi(f) = \text{ess sup } f$ ; or  $\Psi(f) = \int (f'^2/f)$ .

**THEOREM 1.** *Let  $\Psi$  be a functional defined for all densities, and consider a class  $\mathcal{F} = \{f : \Psi(f) \leq c\}$ , where  $c$  is a constant. Assume that there exists an estimate  $\Psi_n = \Psi_n(X_1, \dots, X_n)$  of  $\Psi(f)$  and a sequence  $a_n \rightarrow 0$  such that for all densities  $f$ ,*

$$\sum_{n=1}^{\infty} \mathbf{P} \{ \Psi_n > \Psi(f) + a_n \} < \infty,$$

*and  $\Psi_n \rightarrow \Psi(f)$  almost surely. (Note: the convergence of  $(\Psi_n - \Psi)_+$  must thus be uniform, but not that of  $(\Psi - \Psi_n)_+$ .) Then  $\mathcal{F}$  is testable.*

**REMARK 1.** An explicit test based on Theorem 1 can only be constructed if  $a_n$  is explicitly known. We will provide several examples following the proof.

**PROOF.** Consider the test

$$T_n(X_1, \dots, X_n) = \begin{cases} 1 & \text{if } \Psi_n \leq c + a_n, \\ 0 & \text{otherwise.} \end{cases}$$

If the common density  $f$  of the  $X_i$ 's is in the class  $\mathcal{F}$  then the probability of making a mistake is

$$\mathbf{P} \{ T_n = 0 \} = \mathbf{P} \{ \Psi_n > c + a_n \} \leq \mathbf{P} \{ \Psi_n > \Psi(f) + a_n \},$$

which is summable by assumption, so the Borel-Cantelli lemma implies that the number of mistakes remains finite almost surely. If, on the other hand,  $f \notin \mathcal{F}$ , then  $\Psi(f) > c$ , so since  $\Psi_n \rightarrow \Psi(f)$  almost surely, for sufficiently large  $n$ ,  $\Psi_n > c + a_n$ , and therefore the number of mistakes is finite in this case as well.  $\square$

**EXAMPLE 1: THE CLASS OF DENSITIES SUPPORTED IN  $[-c, c]$ .** As a first simple example, let  $c > 0$ , and consider the class  $\mathcal{F}_c$  of densities supported in the interval  $[-c, c]$ . This class is testable. To prove this, define  $\Psi(f) = \text{ess sup}\{|x| : f(x) > 0\}$ , and  $\Psi_n = \max_{i \leq n} |X_i|$ . The test  $T_n(X_1, \dots, X_n) = I_{\Psi_n \leq c}$  is easily shown to be consistent. However, ignoring the existence of this obvious test, we may also apply Theorem 1. Clearly,  $\Psi_n \rightarrow \Psi(f)$  almost surely. Furthermore, for any  $\epsilon > 0$ ,  $\mathbf{P} \{ \Psi_n > \Psi(f) + \epsilon \} = 0$ , and therefore the condition of Theorem 1 is satisfied for any positive sequence  $a_n \rightarrow 0$ .  $\square$

**EXAMPLE 2: THE CLASS OF DENSITIES BOUNDED BY  $c$ .** Let  $c > 0$ , and consider the class  $\mathcal{F}_c$  of all densities such that  $\text{ess sup } f \leq c$ . Then  $\mathcal{F}_c$  is testable. To prove this statement, we apply Theorem 1 with  $\Psi(f) = \text{ess sup } f$ . The estimate of the functional  $\Psi(f)$  is  $\Psi_n = \Psi(f_n) = \sup_x f_n(x)$ , where  $f_n$  is the kernel estimate of  $f$  defined by

$$f_n(x) = \frac{1}{nh} \sum_{i=1}^n I_{\{|x - X_i| \leq h/2\}},$$

where we choose the smoothing factor  $h$  to be  $h = n^{-1/3}$ . First we check the property involving  $a_n$ . For any  $\epsilon > 0$  and for every  $n$ ,

$$\begin{aligned}
\sup_{f \in \mathcal{F}} \mathbf{P} \{ \Psi_n > \Psi(f) + \epsilon \} &= \sup_{f \in \mathcal{F}} \mathbf{P} \left\{ \sup_x f_n > \text{ess sup } f + \epsilon \right\} \\
&\leq \sup_{f \in \mathcal{F}} \mathbf{P} \left\{ \sup_x \left( \frac{1}{nh} \sum_{i=1}^n I_{\{|x-X_i| \leq h/2\}} - \frac{1}{h} \int_{x-h/2}^{x+h/2} f(z) dz \right) > \epsilon \right\} \\
&\leq \sup_{f \in \mathcal{F}} \left[ \mathbf{P} \left\{ \sup_x \left( \frac{1}{n} \sum_{i=1}^n I_{\{X_i \leq x\}} - \int_{-\infty}^x f(z) dz \right) > \frac{h\epsilon}{2} \right\} \right. \\
&\quad \left. + \mathbf{P} \left\{ \sup_x \left( \int_{-\infty}^x f(z) dz - \frac{1}{n} \sum_{i=1}^n I_{\{X_i \leq x\}} \right) > \frac{h\epsilon}{2} \right\} \right] \\
&\leq 2e^{-nh^2\epsilon^2/2} \\
&= 2e^{-n^{1/3}\epsilon^2/2},
\end{aligned}$$

where the last inequality follows from Massart's (1990) sharpened version of the Dvoretzky-Kiefer-Wolfowitz theorem. Therefore, we may take  $a_n = n^{-1/7}$  in Theorem 1. To complete the proof, we need to show that for any density,

$$\sup_x f_n(x) \rightarrow \text{ess sup } f \quad \text{almost surely.}$$

It is clear from the argument above that for any  $f$ ,  $\limsup_n \Psi_n \leq \Psi(f)$  almost surely. On the other hand, observe that for any  $f$ , by the Lebesgue density theorem (see, e.g., Wheeden and Zygmund, 1977), for every  $B < \text{ess sup } f$  there exists an  $x_0 \in \mathcal{R}$  and a  $\delta_0 > 0$  such that for all  $\delta < \delta_0$ ,  $\int_{x_0-\delta/2}^{x_0+\delta/2} f(z) dz > B\delta$ . Then if  $n$  is so large that  $h = n^{-1/3} < \delta_0$ , then for any  $\epsilon > 0$ ,

$$\begin{aligned}
\mathbf{P} \{ f_n(x_0) < B - \epsilon \} &\leq \mathbf{P} \left\{ \frac{1}{n} \sum_{i=1}^n I_{\{|x_0-X_i| \leq h/2\}} < \int_{x_0-h/2}^{x_0+h/2} f(z) dz - \epsilon h \right\} \\
&\leq e^{-2n\epsilon^2 h^2} = e^{-2n^{1/3}\epsilon^2}
\end{aligned}$$

by Hoeffding's inequality (1963). This, together with the Borel-Cantelli lemma proves that, almost surely,  $\liminf_n \Psi_n \geq \Psi(f)$ , concluding the proof.  $\square$

**EXAMPLE 3: THE CLASS OF DENSITIES WHOSE INTEGRATED SQUARE IS AT MOST  $c$ .** Let  $c > 0$ , and define the class

$$\mathcal{F}_c = \left\{ f : \int f^2 \leq c \right\}.$$

The estimator of  $\Psi(f) = \int f^2$  may be  $\Psi_n = \int f_n^2$  or  $\Psi_n = (2/n) \sum_{i=n/2+1}^n f_{n/2}(X_i)$ , where  $f_n$  is a kernel or histogram estimate of  $f$ . For the sake of simplicity, we take  $\Psi_n = \sum_i N_i^2 / (n^2 h)$ , where  $N_i, i \in \mathcal{Z}$ , are the cardinalities of the intervals  $[ih, ih+h)$ . Observe that changing one  $X_j$  and replacing it by  $X'_j$ , causes at most two  $N_i$ 's to change by one. Therefore, the change in  $\Psi_n$  is at most  $2/(nh)$ . By McDiarmid's inequality (McDiarmid, 1989),

$$\mathbf{P} \{ |\Psi_n - \mathbf{E}\{\Psi_n\}| > \epsilon \} \leq 2 \exp(-nh^2\epsilon^2/2).$$

If  $nh^2/\log n \rightarrow \infty$ , the probability on the left-hand-side is summable, uniformly over all  $f$ . In particular,  $\Psi_n \rightarrow \Psi(f)$  almost surely (even if  $\Psi(f) = \infty$ ) if  $\mathbf{E}\{\Psi_n\} \rightarrow \Psi(f)$ . Denote  $A_i = [hi, h(i+1))$ ,  $p_i = \int_{A_i} f$ .

Then

$$\begin{aligned}
\mathbf{E}\{\Psi_n\} &= \frac{\sum_i (np_i)^2 + np_i(1-p_i)}{n^2h} \\
&= \int g_h^2 + \frac{\sum_i p_i(1-p_i)}{nh} \\
&\quad (\text{where } g_h(x) = p_i/h, x \in A_i \text{ is a density}) \\
&= \int g_h^2 + o(1) .
\end{aligned}$$

Assume  $h \rightarrow 0$  as  $n \rightarrow \infty$ . By the Lebesgue density theorem (Wheeden and Zygmund, 1977),  $g_h \rightarrow f$  at almost all  $x$  when  $h \rightarrow 0$ . Thus, by Fatou's lemma,  $\liminf_{n \rightarrow \infty} \mathbf{E}\{\Psi_n\} \geq \int \liminf_{h \rightarrow 0} g_h^2 = \int f^2$ . This remains valid even if  $\int f^2 = \infty$ . Furthermore, by Jensen's inequality,  $\int g_h^2 \leq \int f^2$ . Collecting all this shows that for all  $f$ ,  $\Psi_n \rightarrow \Psi(f)$  almost surely if  $h \rightarrow 0$ ,  $nh^2/\log n \rightarrow \infty$ . Furthermore, if we take  $a_n = 1/n^{1/5}$ , then for all  $f$ ,

$$\begin{aligned}
\mathbf{P}\{\Psi_n > \Psi(f) + a_n\} &\leq \mathbf{P}\{\Psi_n - \mathbf{E}\{\Psi_n\} > a_n/2\} \quad (\text{if } 1/nh < a_n/2) \\
&\leq 2 \exp(-nh^2/8n^{2/5})
\end{aligned}$$

and this is summable in  $n$  if we take, say,  $h = n^{-1/5}$ . By Theorem 1, the class of densities is thus testable by the test  $\Psi_n \leq c + 1/n^{1/5}$ .  $\square$

EXAMPLE 4: THE CLASS OF DENSITIES  $f$  WITH  $\int f^r \leq c$ , WHERE  $r > 1$  IS A FIXED CONSTANT. The details for this extend those of the previous example, and are left to the reader.  $\square$

### 3. Classes with vanishing minimax risk

Next we derive another general sufficient condition for testability. We recall that a density estimate  $f_n$  is a real-valued measurable function of  $x \in \mathcal{R}$  and the data  $X_1, \dots, X_n$ :

$$f_n(x) = f_n(x, X_1, \dots, X_n).$$

$f_n$  is called **strongly universally consistent** if for every density  $f$ ,

$$\lim_{n \rightarrow \infty} \int |f_n - f| = 0 \quad \text{almost surely.}$$

A density estimate  $f_n$  is **uniformly convergent on  $\mathcal{F}$**  if

$$\lim_{n \rightarrow \infty} \sup_{f \in \mathcal{F}} \mathbf{E} \int |f_n - f| = 0 .$$

Also, we say that  $\mathcal{F}$  is closed (in the  $L_1$  space of densities) if for any density  $g \notin \mathcal{F}$  there exists an  $\epsilon > 0$  and an open ball  $B_{g,\epsilon} = \{f : \int |f - g| < \epsilon\}$  such that  $B_{g,\epsilon} \cap \mathcal{F} = \emptyset$ .

**THEOREM 2.** *Let  $\mathcal{F}$  be a closed class of densities. If there exists a density estimate  $f_n$  that is uniformly convergent on  $\mathcal{F}$ , then  $\mathcal{F}$  is testable. And any closed subclass  $\mathcal{G} \subseteq \mathcal{F}$  is testable.*

In other words, apart from the closedness of the class, we require the convergence to zero of the minimax expected  $L_1$  error of the class, as the following Lemma shows.

LEMMA 1. *The following are equivalent:*

A.

$$\liminf_{n \rightarrow \infty} \sup_{f_n} \sup_{f \in \mathcal{F}} \mathbf{E} \int |f_n - f| = 0 .$$

B. *There exists a density estimate  $f_n$  such that*

$$\lim_{n \rightarrow \infty} \sup_{f \in \mathcal{F}} \mathbf{E} \int |f_n - f| = 0 .$$

PROOF. B implies A. A implies B because A says that for each  $k$ , we may find density estimates  $f_{n,k}$  and numbers  $n_k$  such that for  $n \geq n_k$ ,  $\sup_{f \in \mathcal{F}} \mathbf{E} \int |f_{n,k} - f| < 1/k$ . Without loss of generality,  $n_k \uparrow$ . Define  $g_n = f_{n,k}$ ,  $n_k \leq n < n_{k+1}$ , and use  $g_n$  in part B.  $\square$

It is well-known (Devroye (1983), Birgé (1986)) that very large classes do not satisfy this latter condition. However many positive examples are also known. The theorem also points out that we do not have to explicitly construct  $f_n$ —a proof of existence suffices. Note however that the actual density estimate used to test membership in the class  $\mathcal{F}$  may be very different from the uniformly consistent  $f_n$ . A construction follows from the proof below. The reason we need a special construction is that  $f_n$  may not be stable or concentrated enough to provide good almost sure behavior over an entire sequence.

EXAMPLE 5: FINITE CLASSES OF DENSITIES. As the kernel estimate is universally consistent if we take the smoothing factor  $h$  such that  $h \rightarrow 0$  and  $nh \rightarrow \infty$  ( $nh^d \rightarrow \infty$  in  $\mathbb{R}^d$ ), it is clear from Theorem 2 that we can always test the class  $\mathcal{F} = \{f\}$ . In fact, any finite class of densities is testable.  $\square$

#### 4. Testability via kernel estimates

The next lemma states that to verify the minimax condition, it suffices to establish it for a kernel estimate in which the bandwidth may depend upon the (unknown) density. It removes the burden of construction of minimax optimal or near-optimal densities. Recall that the kernel estimate of a density is

$$f_{nh}(x) = \frac{1}{nh} \sum_{i=1}^n K \left( \frac{x - X_i}{h} \right),$$

where  $h > 0$  is a **bandwidth** and  $K : \mathcal{R} \rightarrow \mathcal{R}$  is a kernel function usually chosen to satisfy  $\int K = 1$ . Note that all kernels used in practice satisfy the condition of the following Lemma.

LEMMA 2. Assume that  $g_n$  is a kernel estimate with kernel  $K$  of polynomial kernel complexity (Devroye and Lugosi, 1997), and with bandwidth  $h_n = h_n(f)$ . Then, if

$$\lim_{n \rightarrow \infty} \sup_{f \in \mathcal{F}} \mathbf{E} \int |g_n - f| = 0,$$

it follows that there exists a density estimate  $f_n$  such that

$$\lim_{n \rightarrow \infty} \sup_{f \in \mathcal{F}} \mathbf{E} \int |f_n - f| = 0,$$

PROOF. This follows from the inequalities of Devroye and Lugosi (1996, 1997), where a data-dependent bandwidth is constructed guaranteeing that the kernel estimate  $f_n$  with that bandwidth satisfies  $\mathbf{E} \int |f_n - f| \leq C \inf_h \mathbf{E} \int |f_{nh} - f| + D \sqrt{\log n/n}$  for universal constants  $D$  and  $C$  not depending upon  $f$ , and all  $n$ . Here  $f_{nh}$  is the kernel estimate with deterministic bandwidth  $h$ .  $\square$

The lemma above allows us to further deduce sufficient conditions for our classes  $\mathcal{F}$ , while avoiding explicit constructions of density estimates for them.

LEMMA 3. If  $\mathcal{G}$  and  $\mathcal{F} \cup \mathcal{G}$  are disjoint classes of densities that are both testable, then so is  $\mathcal{F}$ . If  $\mathcal{F}$  and  $\mathcal{G}$  are testable, then so are  $\mathcal{F} \cup \mathcal{G}$  and  $\mathcal{F} \cap \mathcal{G}$ .

PROOF. We run a test for  $\mathcal{G}$ , which based on a sample of size  $n$  yields the decision  $Y_n$ . Then we run a test on the same sample for membership in  $\mathcal{F} \cup \mathcal{G}$ , and call the decision  $Z_n$ . We decide that the unknown density is in  $\mathcal{F}$  when  $Z_n = 1$  and  $Y_n = 0$ . This makes a finite number of errors with probability one. The second statement does not require disjointness: we decide  $\mathcal{F} \cup \mathcal{G}$  if  $W_n + Y_n \geq 1$ , where  $W_n$  is the decision for  $\mathcal{F}$ . Again, only a finite number of errors are made with probability one. Finally we decide  $\mathcal{F} \cap \mathcal{G}$  if  $W_n Y_n = 1$ .  $\square$

EXAMPLE 6: SCALE/TRANSLATION CLASSES. As a first example, let  $\mathcal{F}$  be any subset of the scale/translation class of densities  $f((\cdot - \mu)/\sigma)/\sigma$ , where  $f$  is a fixed density,  $\mu \in \mathbb{R}$  and  $\sigma > 0$ . Then, in any kernel estimate, take  $h_n = \sigma/\sqrt{n}$ , and note that with this choice, the expected  $L_1$  error is the same for all  $\mu$  and  $\sigma$ . Therefore, by Lemma 2, the  $L_1$  minimax risk tends to zero for this scale/translation class and hence for  $\mathcal{F}$ . Therefore,  $\mathcal{F}$  is testable whenever it is closed. In particular, we can test whether a normal density has a rational  $\sigma = p/q$ , with  $p > 0$  and  $0 < q \leq 500$ . And normality is testable, as is any scale/translation class (by closedness).  $\square$

EXAMPLE 7: NONLINEAR TRANSFORMATION CLASSES. As a second example, let  $\mathcal{F}$  be the class of all densities for random variables  $T(X)$ , where  $X > 0$  has density  $f$ , and  $T(x) = (x^a - 1)/a$ ,  $a > 0$  or  $T(x) = \log(x)$  (case  $a = 0$ ) (the Box-Cox transformations). Then  $\mathcal{F}$  is testable. To see this, transform the data by  $T^{-1}(\cdot)$ , and use a kernel estimate for  $f$  with bandwidth  $h_n = 1/\sqrt{n}$ . Then note that the inequality in the proof of Lemma 2 has been extended to include the joint data-based choice of  $h$  and the Box-Cox parameter  $a$  (Devroye, Lugosi and Udina, 1998). Therefore, using the fact that  $L_1$  errors are invariant under monotone transformations of the input, we conclude there exists a density estimate for which the expected  $L_1$  error tends to zero uniformly over all values of  $a$ . As the density of  $T(X)$  has no limit in the class of densities when  $a \rightarrow \infty$ , we see that  $\mathcal{F}$  is closed, and conclude that  $\mathcal{F}$  is testable.  $\square$

EXAMPLE 8: PARAMETRIC CLASSES. Consider a countable class  $\mathcal{F} = \{\phi_k, 1 \leq k \leq \infty\}$  of densities  $\phi_\infty$  and  $\phi_k$  ( $k < \infty$ ), with  $\phi_k \rightarrow \phi_\infty$ , where convergence is meant in the  $L_1$  sense. This class is closed in the collection of all densities. Also, its  $L_1$  minimax error tends to zero. To see this, consider the kernel estimate with bandwidth  $h_n = 1/\sqrt{n}$ . Then the expected  $L_1$  error is easily bounded as follows: let  $f$  and  $g$  be two densities, and let  $f_n$  and  $g_n$  be two kernel estimates based upon two samples of size  $n$ , both using the same kernel and bandwidth. Then there exists a coupling of the samples such that

$$\mathbf{E} \int |f_n - g_n| \leq \left(1 + \int |K|\right) \int |f - g|$$

(Devroye, 1985). We take  $K$  such that  $\int |K| = 1$ . So, given  $\epsilon > 0$ , let  $K_\epsilon$  be the collection of indices  $k < \infty$  such that  $\int |\phi_k - \phi_\infty| > \epsilon$ . Then, with  $f_{nk}$  denoting the kernel estimate mentioned above (with  $n$  for sample size, and  $k$  for  $\phi_k$ ), we have, letting  $g_n$  denote the kernel estimate (with the same kernel and bandwidth as  $f_{nk}$ ) based on a sample of size  $n$  from  $\phi_\infty$ ,

$$\begin{aligned} & \sup_k \mathbf{E} \int |f_{nk} - \phi_k| \\ & \leq \max_{k \in K_\epsilon} \mathbf{E} \int |f_{nk} - \phi_k| + \sup_{k \notin K_\epsilon} \left( \mathbf{E} \int |f_{nk} - g_n| + \mathbf{E} \int |g_n - \phi| + \int |\phi_k - \phi| \right) \\ & \leq \max_{k \in K_\epsilon} \mathbf{E} \int |f_{nk} - \phi_k| + \mathbf{E} \int |g_n - \phi_\infty| + 3 \sup_{k \notin K_\epsilon} \int |\phi_k - \phi_\infty| \\ & \leq o(1) + 3\epsilon . \end{aligned}$$

Thus,  $\mathcal{F}$  is testable. By arguments as above, the class  $\mathcal{G}$  consisting of all scaled/or translated densities from  $\mathcal{F}$  is also testable. To see why this example is powerful, let  $X_k$  denote a random variable with the gamma density  $x^{k-1}e^{-x}/\Gamma(k)$ ,  $x > 0$ ,  $k > 0$ ,  $k$  integer. Let  $Y_k = (X_k - k)/\sqrt{k}$  be the normalized random variable. Then, as is well-known, the density of  $Y_k$  tends to the normal  $(0, 1)$  density in  $L_1$  as  $k \rightarrow \infty$ . As argued above, the class  $\mathcal{F}$  of all the densities of  $Y_k$ , merged with the standard normal density, is testable. If we consider the scale/translation enlargement  $\mathcal{G}$  of  $\mathcal{F}$ , we obtain all gamma densities, possibly linearly transformed, with integer shape parameter plus all normal densities. It too is testable. Finally, a small additional argument shows that the shape parameter does not have to be restricted at all. Thus, parametric classes in general, when merged with their limit densities, are in most instances testable. When the limits are not included, the question is different, but Lemma 3 provides help. To illustrate this, let  $\mathcal{F}$  be the class of all scaled and translated gamma densities, and  $\mathcal{G}$  be the class of all normal densities. Then, as argued above, both  $\mathcal{F} \cup \mathcal{G}$  and  $\mathcal{G}$  are testable, so  $\mathcal{F}$  is. Lemma 3 is thus useful for the removal of limit classes of parametric collections. Finally, let  $\mathcal{F}$  be the class of all gamma densities, not translated or scaled. This class does not contain any densities as limits, and is thus closed. Hence it is testable.  $\square$



## 5. Testability via total boundedness

A class  $\mathcal{F}$  of densities is said to be totally bounded if for every  $\epsilon > 0$ , there exists a finite number  $N_\epsilon$  of densities  $f_k, 1 \leq k \leq N_\epsilon$  such that the  $L_1$  balls of radius  $\epsilon$  centered at the  $f_k$ 's cover  $\mathcal{F}$ . (Note that in this definition, it is irrelevant whether we additionally ask that the  $f_k$ 's belong to  $\mathcal{F}$ : both definitions would be equivalent.) That is, for every  $f \in \mathcal{F}$ , there exists  $k \leq N_\epsilon$  such that  $\int |f - f_k| \leq \epsilon$ . The smallest possible value of  $\log N_\epsilon$  for  $\mathcal{F}$  is called the Kolmogorov entropy of  $\mathcal{F}$ . Yatracos (1985) (see also Devroye, 1987, p. 90) has constructed a minimum distance estimate  $f_n$  with the property

$$\sup_{f \in \mathcal{F}} \mathbf{E} \left\{ \int |f_n - f| \right\} \leq 5\epsilon + \frac{4 + \sqrt{128N_\epsilon}}{\sqrt{2n}}.$$

Define  $\epsilon_k = 1/k + \inf\{\epsilon > 0 : 128 \log N_\epsilon < \sqrt{k}\}$ . By total boundedness,  $\epsilon_k \rightarrow 0$  as  $k \rightarrow \infty$ . Thus,

$$\sup_{f \in \mathcal{F}} \mathbf{E} \left\{ \int |f_n - f| \right\} \leq 5\epsilon_k + \frac{4 + \sqrt{128N_{\epsilon_k}}}{\sqrt{2n}} \leq o(1) + \frac{4}{\sqrt{2n}} + \frac{8}{n^{1/4}}.$$

Thus, applying Theorem 2, we immediately have

**THEOREM 3.** *Let  $\mathcal{F}$  be a closed totally bounded class of densities. Then  $\mathcal{F}$  is testable.*

**EXAMPLE 9: BOUNDED UNIMODAL DENSITIES.** Let  $\mathcal{F}$  be any closed class of unimodal densities bounded by  $B$  with support on  $[0, 1]$ . Then  $\mathcal{F}$  is testable. In particular, the class of all concave densities with support on  $[0, 1]$  is testable.

**EXAMPLE 10: LIPSCHITZ DENSITIES.** Let  $\mathcal{F}$  be the class of Lipschitz densities on  $[0, 1]$  with Lipschitz constant not exceeding  $C$ . Then  $\mathcal{F}$  or any closed subclass of it is testable.

**EXAMPLE 11: FINITE MIXTURES.** Let  $\mathcal{F}$  be the class of convex mixtures of  $k$  fixed densities. Clearly, this class is closed. Also, by creating a finite grid for the possible convex weights, it is trivial to see that this class is totally bounded. Thus,  $\mathcal{F}$  is testable.

**EXAMPLE 12: UNIFORM MODULUS OF CONTINUITY.** Let  $\mathcal{F}$  be a closed class of densities on  $[0, 1]$  with uniformly bounded modulus of continuity: for all  $\delta > 0$ ,

$$\sup_{f \in \mathcal{F}} \sup_{x, y: \|y-x\| \leq \delta} |f(y) - f(x)| < \infty.$$

Then  $\mathcal{F}$  is totally bounded (Lorentz, 1966; see also Devroye, 1987, p. 98) and thus testable.

## 6. Testability via Yatracos' minimum distance estimates

Next we give another simple sufficient condition for testability via Theorem 2. Recall that given a class of sets  $\mathcal{A}$ , the VC **dimension**  $V$  of  $\mathcal{A}$  is defined as the largest positive integer  $k$  for which there exist  $k$  points  $x_1, \dots, x_k$  such that

$$|\{A \cap \{x_1, \dots, x_k\} : A \in \mathcal{A}\}| = 2^k.$$

If there is no such largest  $k$ , then we say that  $V = \infty$ . If  $V < \infty$ , then  $\mathcal{A}$  is called a VC **class**.

**THEOREM 4.** *Let  $\mathcal{F}$  be a closed class of densities, and assume that the class of sets*

$$\mathcal{A} = \{x : f(x) > g(x); f, g \in \mathcal{F}\}$$

*is a VC class. Then  $\mathcal{F}$  is testable.*

**PROOF.** By Theorem 2, it suffices to prove that there exists a density estimate  $f_n$  which is uniformly convergent in  $\mathcal{F}$ . Consider now the minimum distance estimate proposed by Yatracos (1988):

$$f_n = \arg \min_{f \in \mathcal{F}} \sup_{A \in \mathcal{A}} \left| \int_A f - \mu_n(A) \right|,$$

where  $\mu_n(A) = (1/n) \sum_{i=1}^n I_A(X_i)$  is the empirical measure of  $A$  based on the random sample. Yatracos showed (see also Devroye, Györfi, and Lugosi, 1996, p. 278) that

$$\sup_{f \in \mathcal{F}} \mathbf{E} \int |f_n - f| \leq 4 \mathbf{E} \sup_{A \in \mathcal{A}} \left| \int_A f - \mu_n(A) \right|.$$

The well-known Vapnik-Chervonenkis inequality (Vapnik and Chervonenkis, 1971) implies that

$$\mathbf{E} \left\{ \sup_{A \in \mathcal{A}} \left| \int_A f - \mu_n(A) \right| \right\} \leq 8 \sqrt{\frac{V \log n + 3}{2n}}$$

(see also Devroye, Györfi, and Lugosi, 1996), which finishes the proof.  $\square$

**EXAMPLE 13: EXPONENTIAL MIXTURES.** Let  $k$  be a fixed positive integer, and consider the class  $\mathcal{F}$  of all mixtures of  $k$  exponential densities (that is, translations and scales of  $e^{-x}, x \geq 0$ ). Then Theorem 4 may be applied to show that  $\mathcal{F}$  is testable. It suffices to show that the class

$$\mathcal{A} = \{x : f(x) > g(x); f, g \in \mathcal{F}\}$$

is a VC class. A member set in this class is thus of the form

$$\left\{ x : \sum_{i=1}^{2k} a_i e^{-b_i x} I_{[x > c_i]} > 0 \right\}$$

where  $a_i, c_i \in \mathbb{R}$  and  $b_i > 0$  are free parameters. We claim that each set in this class is the union of at most  $\ell \stackrel{\text{def}}{=} (2k+1)(k+1)$  intervals. Since the class of unions of  $\ell$  intervals is a VC class, we are done. Therefore, the test described above may be used to test whether a density is a  $k$ -mixture of exponentials. We may similarly test for  $k$ -mixtures of normals and indeed many other parametric families. The interval-counting argument is as follows: clearly, we have at most  $k+1$  intervals defined by the thresholds  $c_i$ . It suffices to show that on each of these intervals, a set of the form  $\{x : \sum_{i=1}^{2k} a_i e^{-b_i x} > 0\}$  defines at most  $2k+1$  intervals. But this is well-known (Lemma 25.2 of Devroye, Györfi and Lugosi, 1996). Hence the claim.

## 7. Smoothness and monotonicity classes

In this section, we remove the burden of checking the conditions of Theorem 2, and provide easy-to-verify sufficient conditions for classes of densities to be testable. We begin with a subclass  $\mathcal{F}$  of the monotone densities  $f$  on  $[0, \infty)$  with finite value  $f(0)$ .

**THEOREM 5.** *Let  $\mathcal{F}$  be a closed subclass of the monotone densities  $f$  with finite modal value  $m(f)$ . Assume that*

$$\sup_{f \in \mathcal{F}} \left( \int \sqrt{f} \right)^{\frac{2}{3}} (m(f))^{\frac{1}{3}} < \infty .$$

*Then  $\mathcal{F}$  is testable.*

**PROOF.** Theorem 5 follows from Theorem 2 and Lemmas 1 and 2 if, for  $f \in \mathcal{F}$ , we can establish uniform  $L_1$  error bounds for the kernel estimate  $f_{nh}$  with uniform kernel  $K$  on  $[-1, 1]$ . To this end, we use the typical argument (see Devroye and Györfi, 1985). From the proof below, it will become apparent that the position of the mode is unimportant, and thus we may assume without loss of generality that the mode occurs at zero, and thus that  $m(f) = f(0)$ . Let  $*$  be the convolution operator.

$$\begin{aligned} \mathbf{E} \int |f_{nh} - f| &\leq \int |\mathbf{E}f_{nh} - f| + \int \sqrt{\mathbf{E}(f_{nh} - \mathbf{E}f_{nh})^2} \\ &= \int |f * K_h - f| + \int \sqrt{(1/n)\text{Var}(K_h(x - X_1))} \\ &\leq \int |f * K_h - f| + \int \sqrt{(1/n)\mathbf{E}\{K_h^2(x - X_1)\}} . \end{aligned}$$

For  $x > h$ , observe that  $|f * K_h(x) - f(x)| \leq f(x-h) - f(x+h)$ , and that for  $x < h$ ,  $|f * K_h(x) - f(x)| \leq f(0)$ . Furthermore,  $\mathbf{E}\{K_h^2(x - X_1)\} \leq f(\max(0, x - h))/(2h)$ . Using these estimates, we obtain

$$\begin{aligned} \mathbf{E} \int |f_{nh} - f| &\leq hf(0) + \int_h^{3h} f + h\sqrt{\frac{f(0)}{2nh}} + \frac{\int \sqrt{f}}{\sqrt{2nh}} \\ &\leq 3hf(0) + h\sqrt{\frac{f(0)}{2nh}} + \frac{\int \sqrt{f}}{\sqrt{2nh}} \\ &= \frac{4(\int \sqrt{f})^{\frac{2}{3}}(f(0))^{\frac{1}{3}}}{(2n)^{\frac{1}{3}}} + \frac{(\int \sqrt{f})^{\frac{1}{3}}(f(0))^{\frac{1}{6}}}{(2n)^{\frac{2}{3}}} \end{aligned}$$

when we take  $h^{3/2} = \int \sqrt{f}/(f(0)\sqrt{2n})$ . Therefore, we have uniform convergence to zero whenever

$$\sup_{f \in \mathcal{F}} \left( \int \sqrt{f} \right)^{\frac{2}{3}} (f(0))^{\frac{1}{3}} < \infty .$$

By Lemma 2 and Theorem 2, Theorem 5 follows.  $\square$

Let us extend the previous Theorem to include densities of bounded variation. We recall that a density  $f$  is of bounded variation if we may find an increasing function  $f_1$  and a decreasing function  $f_2$  such that  $f = f_1 + f_2$ , such that the total variation

$$\mathcal{V}(f) \stackrel{\text{def}}{=} \inf_{f_1, f_2: f=f_1+f_2, f_1 \uparrow, f_2 \downarrow} \sup_{y > x} (f_1(y) - f_1(x)) + \sup_{y > x} (f_2(x) - f_2(y)) < \infty .$$

THEOREM 6. Let  $\mathcal{F}$  be a closed subclass of the densities of bounded variation, and assume that

$$\sup_{f \in \mathcal{F}} (\mathcal{V}(f))^{\frac{1}{3}} \left( \int \sqrt{f} \right)^{\frac{2}{3}} < \infty .$$

Then  $\mathcal{F}$  is testable.

PROOF. We argue as in the proof of Theorem 5, and apply Lemma 2 and Theorem 2. Let  $f_{nh}$  be the kernel estimate. If  $f = f_1 + f_2$  is the bounded variation decomposition of  $f$ , with  $f_1 \uparrow$  and  $f_2 \downarrow$ , and  $K$  is the uniform kernel on  $[-1, 1]$ , then

$$|f * K_h - f| \leq f_1(x+h) - f_1(x) + f_2(x-h) - f_2(x)$$

and

$$|f * K_h - f| \leq 2h\mathcal{V}(f) .$$

Also,

$$\int \sqrt{\frac{\mathbf{E}\{K_h^2(x - X_1)\}}{n}} \leq \int \sqrt{\frac{f(x+h) + f(x-h)}{2hn}} dx \leq \sqrt{\frac{2}{nh}} \int \sqrt{f}$$

so that

$$\mathbf{E} \{|f_{nh} - f|\} \leq 2h\mathcal{V}(f) + \sqrt{\frac{2}{nh}} \int \sqrt{f} = \frac{4}{(2n)^{1/3}} \left( \int \sqrt{f} \right)^{\frac{2}{3}} (\mathcal{V}(f))^{\frac{1}{3}}$$

if we take  $h^{3/2} = \int \sqrt{f} / (\sqrt{2n}\mathcal{V}(f))$ .  $\square$

EXAMPLE 14: LOG-CONCAVE DENSITIES. A very important subclass of the densities is the class of log-concave densities:  $\log f$  is concave. We claim that any closed subclass of the log-concave densities is testable. Consider that this class includes all beta densities with both parameters greater than or equal to 1, all gamma densities with shape parameter greater than or equal to one, all exponential power distributions with shape parameter at least one, the normal densities, and a host of other densities. It is known that these densities are unimodal and that if the mode occurs at  $z$ , a rescaling of the random variable to place the mode at zero with modal value one results in a density  $g$  with  $g(u) \leq e^{1-|u|}$  (Devroye, 1987). As the product in Theorem 5 is scale and translation invariant, and the log-concave inequality is absolute, we see that, except for the monotonicity, all conditions of Theorem 5 are fulfilled. It is left as a trivial exercise to extend Theorem 5 to this class of densities. Thus, we have a very simple condition for establishing the testability of large subclasses of the famous parametric families. And as the class of log-concave densities is closed, we can indeed test log-concavity.  $\square$

EXAMPLE 15: CONCAVE DENSITIES. Consider the class of all concave densities on their support. Clearly, this class is closed. Furthermore, if the mode is forced to be at zero and of modal value one, then any density in this class is bounded by one, and of support contained in  $[-2, 2]$ . By an argument as for the log-concave densities, this class is testable (regardless of the support!), and indeed any closed subclass of it is testable as well.  $\square$

One could refine the bounds of the proof of Theorem 5 and indeed use higher-order kernels to obtain results for subclasses related to Akhiezer classes of densities. In this manner, we may deal with convex subclasses of the monotone densities, and classes for which the  $r$ -th derivative  $f^{(r)}$  is monotone.

EXAMPLE 16: LIPSCHITZ DENSITIES. A class  $\mathcal{F}$  of Lipschitz (1) densities with Lipschitz constant  $C(f) < \infty$  and support  $s(f)$  is testable if  $\mathcal{F}$  is closed and

$$\sup_{f \in \mathcal{F}} C(f)s^2(f) < \infty .$$

To see this, apply Theorem 6, and note that  $\mathcal{V}(f) \leq C(f)s(f)$ , and  $\int \sqrt{f} \leq \sqrt{s(f) \int f} = \sqrt{s(f)}$ .  $\square$

EXAMPLE 17: UNIMODAL DENSITIES. Let  $\mathcal{F}$  be a class of unimodal densities with modal value  $m(f)$  and variance  $\sigma^2(f)$ . Then this class is testable if  $\mathcal{F}$  is closed and

$$\sup_{f \in \mathcal{F}} m(f)(1 + \sigma^2(f)) < \infty .$$

To see this, note that  $\mathcal{V}(f) \leq 2m(f)$  and assuming without loss of generality that the mean is at the origin,

$$\int \sqrt{f} = \int \frac{\sqrt{(1+x^2)f}}{\sqrt{1+x^2}} \leq \sqrt{\int (1+x^2)f} \sqrt{\int \frac{1}{1+x^2}} = \sqrt{\pi(1+\sigma^2(f))} .$$

Examples of such classes include the gamma densities with shape parameter 1 or larger, the symmetric beta densities with shape parameter 1 or larger, and the unimodal densities  $f \leq A/(1+x^4)$ , with  $A < \infty$  fixed and given.  $\square$

## 8. Ad hoc analysis: testing unimodality and convexity

Surprisingly, there are classes whose minimax risk and whose complement's minimax risk does not tend to zero, and yet they are testable. We have already encountered such classes in Section 2. In this section we construct explicit tests to prove that the following classes are testable: the monotone densities on  $[0, \infty)$ , the monotone densities on an interval of the real line, the unimodal densities with mode at 0, and the unimodal densities. To keep the material limited, we will only provide an explicit proof for the class of monotone densities on  $[0, \infty)$ . We recall from Devroye (1983) that the minimax risk of this class does not tend to zero. To construct an explicit ad hoc test, consider a histogram estimate with bins  $[0, h), [h, 2h), [2h, 3h), \dots$ . Let  $N_i$  denote the number of data points in  $[ih, (i+1)h)$ . The test is the following: take  $h = 1/n^{1/7}$ . Then

$$\text{decide non-monotone if } \max_{i \geq 0} (N_{i+1} - N_i) > n^{2/3} ;$$

decide monotone otherwise.

THEOREM 7. *The test above makes almost surely a finite number of errors for testing membership in the class of monotonically decreasing densities on  $[0, \infty)$ .*

PROOF. First, assume that  $f$  is indeed monotone on  $[0, \infty)$ . We bound  $\mathbf{P}\{N_{i+1} - N_i > t\}$  by standard methods: For each  $i \geq 0$ , define  $p_{ni} = \mathbf{P}\{X_1 \in [ih, (i+1)h)\}$ . Introduce  $N'_i$  such that  $(N'_i, N_i)$  are two components of a multinomial random vector with total count  $n$  and success probabilities  $p_{ni}$  each. Then

$$\begin{aligned} \mathbf{P}\{N_{i+1} - N_i > t\} &\leq \mathbf{P}\{N'_i - N_i > t\} \\ &\leq \mathbf{P}\{N'_i - np_{ni} > t/2\} + \mathbf{P}\{N_i - np_{ni} < -t/2\} \\ &= \mathbf{P}\{|N_i - np_{ni}| > t/2\} \\ &\leq 2e^{-\frac{t^2}{2n}}, \end{aligned}$$

by Hoeffding's inequality (Hoeffding, 1963). Thus,

$$\begin{aligned} &\mathbf{P}\left\{\max_{i \geq 0}(N_{i+1} - N_i) > n^{2/3}\right\} \\ &\leq \mathbf{P}\left\{\max_{i \geq n} N_i > n^{2/3}\right\} + 2ne^{-n^{1/3}/2} \\ &\leq \sum_{i \geq n} \mathbf{P}\left\{N_i > n^{2/3}\right\} + 2ne^{-n^{1/3}/2} \\ &\leq \sum_{i \geq n} \mathbf{P}\left\{\text{binomial}(n, 1/(i+1)) > n^{2/3}\right\} + 2ne^{-n^{1/3}/2} \\ &\quad (\text{because } p_{ni} \leq 1/(i+1) \text{ by monotonicity}) \\ &\leq \sum_{i \geq n} \left(\frac{n}{(i+1)n^{2/3}}\right)^{n^{2/3}} \cdot \frac{i+1}{i} + 2ne^{-n^{1/3}/2} \\ &\quad (\text{because } \mathbf{P}\{\text{binomial}(n, p) \geq t\} \leq \binom{n}{t} p^t / (1-p) \leq (np/t)^t / (1-p), t \text{ integer}) \\ &= O\left(e^{-n^{1/3}/3}\right). \end{aligned}$$

In the above chain, we assumed without loss of generality that  $n^{2/3}$  is integer-valued. As this is summable in  $n$ , the Borel-Cantelli lemma implies that we will make finitely many errors almost surely.

Next assume that  $f$  is not monotone. Thus, there exist Lebesgue points  $0 < y < z$  with the property that  $f(y) = f(z) - 3\delta$  for some  $\delta > 0$ . As these are Lebesgue points, we know that there exists an  $\epsilon > 0$  such that for any interval  $I$  of length  $|I| < \epsilon$  containing  $y$ ,  $\int_I f(x)dx < |I|(f(y) + \delta)$ , and similarly for  $z$ :  $\int_I f(x)dx > |I|(f(z) - \delta)$ . The number of intervals  $[ih, (i+1)h)$  separating  $z$  from  $y$  is at most  $2 + (z - y)/h$ . Let  $h < \epsilon$ , and let  $i$  range over the intervals covering  $[y, z]$ . The difference between the last  $p_{ni}$  and the first  $p_{ni}$  is at least  $\delta h$ , so that the maximal differential  $p_{n,i+1} - p_{n,i}$  for  $i$  and  $i+1$  both among the given intervals is at least  $\delta h / (2 + (z - y)/h) \geq \delta h^2 / 2(z - y) \stackrel{\text{def}}{=} 3ch^2$  for  $n$  large enough (and thus  $h$  small enough). Let  $j$  be an index for which  $p_{n,j+1} - p_{n,j} \geq 3ch^2$ . If we decide that  $f$  is monotone, then we have  $N_{j+1} - N_j \leq n^{2/3}$ , however. Thus, the probability of erring is not more than

$$\begin{aligned} \mathbf{P}\{N_{j+1} - N_j \leq n^{2/3}\} &\leq \mathbf{P}\{N_{j+1} - np_{n,j+1} - (N_j - np_{n,j}) \leq n^{2/3} - 3cnh^2\} \\ &= \mathbf{P}\{N_{j+1} - np_{n,j+1} - (N_j - np_{n,j}) \leq n^{2/3} - 3cn^{5/7}\} \\ &\leq \mathbf{P}\{N_{j+1} - np_{n,j+1} - (N_j - np_{n,j}) \leq -2cn^{5/7}\} \end{aligned}$$

$$\begin{aligned}
& \text{(for } n \text{ large enough)} \\
& \leq \mathbf{P}\{N_{j+1} - np_{n,j+1} < -cn^{5/7}\} + \mathbf{P}\{N_j - np_{nj} > cn^{5/7}\} \\
& \leq 2e^{-2c^2 n^{3/7}}
\end{aligned}$$

by Hoeffding's inequality. By the Borel-Cantelli lemma, we once again make finitely many errors almost surely.  $\square$

For monotonicity on any right-infinite interval, we apply the same test, starting with the second occupied interval from the left. For unimodality, we find the interval of maximal cardinality, and apply monotonicity tests of opposite polarity on both sides of that maximal interval. The details are uninteresting. It is also worth noting that we can test for convexity. In particular, if  $i$  and  $j$  are the indices of the leftmost and rightmost intervals in a grid of intervals of width  $h$  that are occupied, then the test with  $h = 1/n^{1/7}$  that decides against convexity if  $\max_{i < k < j-2} (2N_{k+1} - N_{k+2} - N_k) > n^{2/3}$  errs finitely often with probability one, for any density. In fact, one can in this manner test membership in any Akhiezer class.

## 9. Proof of Theorem 2

The proof of the theorem is split into two lemmas, the first of which is directly applicable to prove testability in many cases.

LEMMA 4. *Let  $\mathcal{F}$  be a closed class of densities. If there exists a sequence of positive numbers  $b_n \rightarrow 0$  and a density estimate  $g_n$  such that  $g_n$  is strongly universally consistent, moreover for every  $f \in \mathcal{F}$*

$$\sum_{n=1}^{\infty} \mathbf{P} \left\{ \int |g_n - f| > b_n \right\} < \infty,$$

*then  $\mathcal{F}$  is testable.*

PROOF. Consider the test

$$T_n(X_1, \dots, X_n) = \begin{cases} 1 & \text{if } \inf_{f \in \mathcal{F}} \int |g_n - f| \leq b_n, \\ 0 & \text{otherwise.} \end{cases}$$

In other words, based on the data  $X_1, \dots, X_n$ , we compute the density estimate  $g_n$ , and project it in the  $L_1$  distance on the class  $\mathcal{F}$ . If the distance between  $g_n$  and the class is greater than  $b_n$ , then we say that the unknown density is not in the class. Now it is easy to see that this test is consistent, since if the  $X_i$ 's are drawn from a density  $f \in \mathcal{F}$  then

$$\mathbf{P}\{T_n \text{ fails}\} = \mathbf{P} \left\{ \inf_{g \in \mathcal{F}} \int |g_n - g| > b_n \right\} \leq \mathbf{P} \left\{ \int |g_n - f| > b_n \right\}.$$

By assumption these probabilities are summable over  $n = 1, 2, \dots$ , so the Borel-Cantelli lemma implies that, almost surely,  $T_n$  fails at most for finitely many  $n$ .

On the other hand, if  $f \notin \mathcal{F}$ , then by closedness of the class  $\mathcal{F}$ , there exists an  $\epsilon > 0$  such that  $\inf_{g \in \mathcal{F}} \int |f - g| > \epsilon$ . However, since  $g_n$  is strongly universally consistent and  $b_n \rightarrow 0$ , eventually, almost surely,  $\inf_{f \in \mathcal{F}} \int |g_n - f| > \epsilon/2 > b_n$ , and therefore the test does not fail.  $\square$

In many cases there exists a simple estimate which satisfies the condition of Lemma 4. For example, if  $\mathcal{F}$  is the class of all densities supported on  $[a, b]$  such that  $\text{ess sup}_{x,y} |f(x) - f(y)|/|x - y| \leq c$ , then the kernel estimate with an appropriate non-data-dependent bandwidth will do. However, even simple cases as the class of all normal densities, any such kernel estimate fails to provide a uniform rate of convergence within  $\mathcal{F}$ . In this specific case it is possible to define a data-dependent bandwidth with the desired property (see Devroye, 1989). However, to finish the proof of Theorem 2, we need a universal construction, provided in the lemma below:

LEMMA 5. *If for some sequence  $a_n \rightarrow 0$  there exists a density estimate  $f_n$  with  $\sup_{f \in \mathcal{F}} \mathbf{E} \int |f_n - f| \leq a_n$ , then there exists another sequence  $b_n \rightarrow 0$  and a density estimate  $g_n$  such that  $g_n$  is strongly universally consistent, and for all  $f \in \mathcal{F}$*

$$\sum_{n=1}^{\infty} \mathbf{P} \left\{ \int |g_n - f| > b_n \right\} < \infty.$$

REMARK 2. The proof below shows that one may always take  $b_n = 3a_{\lfloor (n/2)^{1/3} \rfloor} + 3(n/2)^{-1/12} + 4n^{-1/3}$ , so  $b_n$  only depends on  $a_n$  but not on the class  $\mathcal{F}$ . However, in most cases this is a suboptimal choice. The estimate  $g_n$  defined below is merely a part of a general proof of existence. In most concrete cases much superior estimates exist. Also, in fact, we show that  $\sup_{f \in \mathcal{F}} \sum_{n=1}^{\infty} \mathbf{P} \{ \int |g_n - f| > b_n \} < \infty$ .

PROOF. First, we “stabilize”  $f_n$  to make sure that the  $L_1$  error  $\int |f_n - f|$  is always concentrated around its mean, and then combine it with a consistent estimate to achieve strong universal consistency. We define the stabilized density estimate  $\hat{f}_n$  as follows:

$$\hat{f}_n(x, X_1, \dots, X_n) = \frac{1}{\lfloor \frac{n}{N} \rfloor} \sum_{k=1}^{\lfloor \frac{n}{N} \rfloor} f_N(x, X_{(k-1)N+1}, \dots, X_{kN}),$$

where  $N = \lfloor n^{1/3} \rfloor$ . In other words, we chop up the data into about  $n^{2/3}$  equal blocks, construct the estimate on all blocks, and take their average. McDiarmid’s inequality (1989) assures that if by changing the value of one data point but leaving all others intact the  $L_1$  error does not change by much, then it is close to its mean with large probability. In our case,

$$\begin{aligned} & \left| \int |\hat{f}_n(x, X_1, \dots, X_n) - f(x)| dx - \int |\hat{f}_n(x, X_1, \dots, X'_i, \dots, X_n) - f(x)| dx \right| \\ & \leq \left| \int \hat{f}_n(x, X_1, \dots, X_n) - \int \hat{f}_n(x, X_1, \dots, X'_i, \dots, X_n) dx \right| \\ & \leq \frac{1}{\lfloor \frac{n}{N} \rfloor} \int |f_N(x, X_{(k-1)N+1}, \dots, X_{kN}) - f_N(x, X_{(k-1)N+1}, \dots, X'_i, \dots, X_{kN})| dx \\ & \quad \text{(if the index } i \text{ is in the } k\text{-th block)} \\ & \leq \frac{2}{\lfloor \frac{n}{N} \rfloor}, \end{aligned}$$

where the last inequality follows from the fact that we may assume that the estimate  $f_N$  is always a density (i.e., nonnegative and integrates to one, since otherwise with standard operations one may always



construct such an estimate by not increasing the  $L_1$  error, see Devroye (1987)), and that the  $L_1$  distance between any two densities is at most 2. Thus, McDiarmid's inequality implies that for any  $t > 0$ ,

$$\mathbf{P} \left\{ \int |\widehat{f}_n - f| - \mathbf{E} \int |\widehat{f}_n - f| > t \right\} \leq e^{-t^2/2n(1/\lfloor \frac{n}{8} \rfloor)^2} \leq e^{-t^2 n^{1/3}/8}.$$

if  $n \geq 3$ . Thus, for example, by taking  $t = n^{-1/12}$  and using the simplified notation  $a'_n = a_{\lfloor n^{1/3} \rfloor} + n^{-1/12}$ , for any  $f \in \mathcal{F}$ ,

$$\mathbf{P} \left\{ \int |\widehat{f}_n - f| > a'_n \right\} \leq e^{-n^{1/6}/8}.$$

The second step is to extend  $\widehat{f}_n$  so that it becomes universally consistent. Without loss of generality we may assume again that  $\widehat{f}_n$  is indeed a density. Let  $\xi_n$  be an arbitrary strongly universally consistent density estimate. We proceed as follows: Split the available data into two equal parts  $X_1, \dots, X_{n/2}$  and  $X_{n/2+1}, \dots, X_n$ . Based on the first half construct the estimates  $\widehat{f}_{n/2}$  and  $h_{n/2}$ . Define the ‘‘Yatracos set’’ (see Yatracos (1985) or Devroye and Lugosi (1997))

$$A_{n/2} = \left\{ x : \widehat{f}_{n/2}(x) > \xi_{n/2}(x) \right\},$$

and use the second half of the data to calculate the empirical probability

$$\mu_{n/2}(A_{n/2}) = \frac{2}{n} \sum_{j=n/2+1}^n I_{\{X_j \in A_{n/2}\}}.$$

The density estimate  $g_n$  is defined by

$$g_n = \begin{cases} \widehat{f}_{n/2} & \text{if } \left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} \widehat{f}_{n/2} \right| < \left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} \xi_{n/2} \right|; \\ \xi_n & \text{otherwise.} \end{cases}$$

We need to show that  $g_n$  is universally consistent, and within  $\mathcal{F}$  it has a uniform rate of convergence. We start with the case  $f \in \mathcal{F}$ . Define  $c_n = 2a'_{n/2} + 4n^{-1/3}$ . If  $\int |\widehat{f}_{n/2} - \xi_{n/2}| \leq c_n$  then by the triangle inequality

$$\int |g_n - f| \leq \int |\widehat{f}_{n/2} - f| + c_n.$$

Otherwise, if  $\int |\widehat{f}_{n/2} - \xi_{n/2}| = 2 \left( \int_{A_{n/2}} \widehat{f}_{n/2} - \int_{A_{n/2}} \xi_{n/2} \right) > c_n$ , observing that

$$\left| \int_{A_{n/2}} f - \int_{A_{n/2}} \widehat{f}_{n/2} \right| \leq \frac{1}{2} \int |\widehat{f}_{n/2} - f|$$

by Scheffé's theorem, we have  $g_n = \widehat{f}_{n/2}$  whenever

$$\left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} f \right| < \frac{c_n}{4} - \frac{1}{2} \int |\widehat{f}_{n/2} - f|.$$

Summarizing the two cases, we see that for all  $f \in \mathcal{F}$ ,

$$\begin{aligned} \mathbf{P} \left\{ \int |g_n - f| > a'_{n/2} + c_n \right\} &\leq \mathbf{P} \left\{ \int |g_n - f| > \int |\widehat{f}_{n/2} - f| + c_n \right\} + e^{-(n/2)^{1/6}/8} \\ &\leq \mathbf{P} \left\{ \left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} f \right| > \frac{c_n}{4} - \frac{a'_{n/2}}{2} \right\} + 2e^{-(n/2)^{1/6}/8} \\ &\quad \text{(by the above argument)} \end{aligned}$$

$$\begin{aligned}
&= \mathbf{P} \left\{ \left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} f \right| > n^{-1/3} \right\} + 2e^{-(n/2)^{1/6}/8} \\
&\leq e^{-n^{1/3}} + 2e^{-(n/2)^{1/6}/8} \\
&\quad (\text{by Hoeffding's inequality, 1963}).
\end{aligned}$$

Therefore, taking  $b_n = a'_{n/2} + c_n$ , we have that, indeed, for every  $f \in \mathcal{F}$ ,

$$\sum_{n=1}^{\infty} \mathbf{P} \left\{ \int |g_n - f| > b_n \right\} < \infty.$$

Now it remains to show that  $g_n$  is strongly universally consistent. This may be done in a similar way: If  $\int |\widehat{f}_{n/2} - \xi_{n/2}| \leq 2 (\int |\xi_{n/2} - f| + n^{-1/3})$  then by the triangle inequality,

$$\int |g_n - f| \leq 3 \int |\xi_{n/2} - f| + 2n^{-1/3}.$$

Otherwise, by Scheffé's theorem,  $\int_{A_{n/2}} \widehat{f}_{n/2} - \int_{A_{n/2}} \xi_{n/2} > \int |\xi_{n/2} - f| + n^{-1/3}$  and

$$\left| \int_{A_{n/2}} f - \int_{A_{n/2}} \xi_{n/2} \right| \leq \frac{1}{2} \int |\xi_{n/2} - f|.$$

Therefore,  $g_n = \xi_{n/2}$  whenever

$$\left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} f \right| < \frac{1}{2} n^{-1/3}.$$

Therefore, in all cases, since the  $L_1$  distance is bounded by 2,

$$\int |g_n - f| \leq 3 \int |\xi_{n/2} - f| + 2n^{-1/3} + 2I \left\{ \left| \mu_{n/2}(A_{n/2}) - \int_{A_{n/2}} f \right| \geq (1/2)n^{-1/3} \right\}.$$

By the strong universal consistency of  $\xi_n$  and Hoeffding's inequality, all terms on the right-hand side converge to zero almost surely. This concludes the proof of Lemma 5 and Theorem 2.  $\square$

## 10. Non-testable classes

In this section we establish sufficient conditions for the nontestability of a class of densities, and show several examples of nontestable classes.

We first generalize the definition of testability: if  $\mathcal{X}$  is a class of densities, then we say that  $\mathcal{F}$  is **testable with respect to**  $\mathcal{X}$  if there exists a consistent test to decide whether  $f \in \mathcal{F} \cap \mathcal{X}$  or  $f \in \mathcal{F}^c \cap \mathcal{X}$ . Clearly, if  $\mathcal{F}$  is not testable with respect to  $\mathcal{X}$ , then  $\mathcal{F}$  is not testable (with respect to the class of all densities).

It will be convenient to work, instead of densities, with the inverse of their corresponding cumulative distribution function: Any density  $f$  is uniquely determined by a monotonically increasing function  $G : (0, 1) \rightarrow (-\infty, \infty)$  defined by  $G = F^{-1}$ , where  $F(x) = \int_{-\infty}^x f(z) dz$ . If  $U_1, U_2, \dots$  is a sequence of independent uniform  $[0, 1]$  random variables, then  $G(U_1), G(U_2), \dots$  is a sequence of i.i.d. random variables with density  $f$ . This is the coupling between samples from different distributions which will be used in the proof below.

THEOREM 8. Let  $\mathcal{F}, \mathcal{X}$  be classes of densities. Denote the class of all inverse distribution functions corresponding to densities in  $\mathcal{X}$  by  $\mathcal{A}$ , and to those in  $\mathcal{F} \cap \mathcal{X}$  by  $\mathcal{G}$ . Assume that there exist two sets of functions  $\mathcal{B} \subset \mathcal{G}$  and  $\mathcal{C} \subset \mathcal{A} - \mathcal{G}$  with the following property:

- (1) there exists a family of subsets  $S_\epsilon$  of  $(0, 1)$  indexed by real numbers  $\epsilon \in (0, 1)$  such that  $\lambda(S_\epsilon) \leq \epsilon$  ( $\lambda$  denotes the Lebesgue measure);
- (2) if  $\epsilon_1 > \epsilon_2$  then  $S_{\epsilon_1} \supseteq S_{\epsilon_2}$  ;
- (3) for any  $\epsilon \in (0, 1)$  and  $G \in \mathcal{B}$  there exists a  $H \in \mathcal{C}$  such that  $G(x) = H(x)$  for all  $x \notin S_\epsilon$ .
- (4) for any  $\epsilon \in (0, 1)$  and  $H \in \mathcal{C}$  there exists a  $G \in \mathcal{B}$  such that  $G(x) = H(x)$  for all  $x \notin S_\epsilon$ .

Then  $\mathcal{F}$  is not testable with respect to  $\mathcal{X}$ .

PROOF. Let  $U_1, U_2, \dots$  be an i.i.d. sequence of uniform  $[0, 1]$  random variables, from which we obtain all samples for all distributions by the inverse distribution function transformation. We may represent this sequence by the probability element  $\omega$ . Assume that there exists a consistent test  $T_n$ . Then for any density  $f \in \mathcal{F} \cap \mathcal{X}$ , and almost all  $\omega$  (i.e., with probability one) there exists an integer  $N(\omega)$  such that

$$T_n(X_1, \dots, X_n) = 1 \quad \text{if } n > N(\omega)$$

and for any density  $f \in \mathcal{F}^c \cap \mathcal{X}$ , and almost all  $\omega$  there exists an integer  $N(\omega)$  such that

$$T_n(X_1, \dots, X_n) = 0 \quad \text{if } n > N(\omega).$$

We will construct a density such that, with probability more than  $1/2$ ,  $T_n(X_1, \dots, X_n) = 0$  for infinitely many  $n$  and  $T_n(X_1, \dots, X_n) = 1$  for infinitely many  $n$ , which is a contradiction.

We use the coupling defined in the introduction of this section. Let  $\delta_k = 2^{-k-2}$ ,  $k = 1, 2, \dots$ , and let  $G_1 \in \mathcal{B}$  be arbitrary. Then there exists an integer  $N_1 = N_1(\omega)$  such that

$$\mathbf{P} \{T_n(G_1(U_1), \dots, G_1(U_n)) = 1 \text{ for all } n \geq N_1\} > 1 - \delta_1$$

(see, e.g., Royden, 1968, p. 70, Problem 23.a). Choose  $\epsilon_1 > 0$  such that  $(1 - \epsilon_1)^{N_1} > 1 - \delta_1$ , and consider a function  $G_2 \in \mathcal{C}$  which agrees with  $G_1$  on  $(0, 1) - S_{\epsilon_1}$ . Then

$$\mathbf{P} \{G_1(U_1) = G_2(U_1), \dots, G_1(U_{N_1}) = G_2(U_{N_1})\} > 1 - \delta_1,$$

and

$$\mathbf{P} \{T_n(G_2(U_1), \dots, G_2(U_{N_1})) = 1\} > 1 - 2\delta_1.$$

Now similarly, since  $G_2 \in \mathcal{C}$ , there exists an integer  $N_2 = N_2(\omega) > N_1$  such that

$$\mathbf{P} \{T_n(G_2(U_1), \dots, G_2(U_n)) = 0 \text{ for all } n \geq N_2\} > 1 - \delta_2.$$

Next we choose  $\epsilon_2 > 0$  such that  $(1 - \epsilon_2)^{N_2} > 1 - \delta_2$ , and consider a function  $G_3 \in \mathcal{B}$  which agrees with  $G_2$  on  $(0, 1) - S_{\epsilon_2}$ . We continue this procedure such that  $G_k \in \mathcal{B}$  for odd  $k$  and  $G_k \in \mathcal{C}$  for even  $k$ , and  $G_k$  agrees with  $G_{k-1}$  on  $(0, 1) - S_{\epsilon_{k-1}}$ , where  $(1 - \epsilon_{k-1})^{N_{k-1}} > 1 - \delta_{k-1}$ , and  $N_k$  is chosen such that

$$\mathbf{P} \left\{ T_n(G_k(U_1), \dots, G_k(U_n)) = I_{\{k \text{ is odd}\}} \text{ for all } n \geq N_k \right\} > 1 - \delta_k.$$

Then clearly the sequence of these functions  $G_1, G_2, \dots$  converges pointwise to some function  $G \in \mathcal{A}$ . Also,

$$\mathbf{P} \left\{ T_n(G(U_1), \dots, G(U_{N_k})) = I_{\{k \text{ is odd}\}} \text{ for all } k = 1, 2, \dots \right\} > 1 - \sum_{k=1}^{\infty} 2\delta_k = \frac{1}{2},$$

and the proof is finished.  $\square$

EXAMPLE 18: BOUNDEDNESS OF THE SUPPORT OF A DENSITY IS NOT TESTABLE. Let  $\mathcal{F}$  be the class of all densities whose support is bounded. Then Theorem 8 implies that  $\mathcal{F}$  is not testable. To see this, simply take  $S_\epsilon = (1 - \epsilon, 1)$ , and let  $\mathcal{B}$  be the family of inverse distribution functions corresponding to positive bounded random variables with a density, and let  $\mathcal{C}$  be the family corresponding to positive unbounded random variables with a density. Then clearly, functions in  $\mathcal{B}$  are bounded and functions in  $\mathcal{C}$  are unbounded, and for any bounded function  $G \in \mathcal{B}$  and  $\epsilon \in (0, 1)$  there exists an unbounded  $H \in \mathcal{C}$  such that  $G$  and  $H$  agree on  $(0, 1 - \epsilon)$  and vice versa, and therefore the condition of Theorem 8 is satisfied. It is also easy to see that one cannot test boundedness of the support with respect to the following classes: all bounded densities, all continuous densities, all Lipschitz densities, and all unimodal densities. In particular, unimodality is testable, but not bounded support once it is known that a density is unimodal.

EXAMPLE 19: BOUNDEDNESS OF A DENSITY IS NOT TESTABLE. Let  $\mathcal{F}$  be the class of all densities with  $\text{ess sup } f < \infty$ . Then  $\mathcal{F}$  is not testable. We show this, via Theorem 8, by proving that  $\mathcal{F}$  is not testable with respect to  $\mathcal{X}$ , the class of all monotonically decreasing densities on  $[0, \infty)$  that are continuous on  $(0, \infty)$ . Then all inverse distribution functions  $G \in \mathcal{A}$  corresponding to densities in  $\mathcal{X}$  are concave increasing functions with  $G(0) = 0$ ,  $G'(0) \geq 0$ , and  $\liminf_{t \downarrow 0} G'(t) = \infty$  if and only if the density is unbounded. Take  $S_\epsilon = (0, \epsilon)$ , and continue a function  $G$  for an unbounded density on  $[0, \epsilon]$  by a parabola through  $(0, 0)$  and  $(\epsilon, G(\epsilon))$  with derivative  $G'(\epsilon)$  at  $\epsilon$  (which corresponds to a  $G$  for a bounded continuous and monotonically decreasing density on  $[0, \infty)$ ). Similarly, continue a function  $G$  for a bounded density on  $[0, \epsilon]$  by a quadratic Bezier spline (Farin, 1993) having derivative  $\infty$  at 0,  $G'(\epsilon)$  at  $\epsilon$ , and with control point at  $(0, a)$ , where  $a$  is the place where the line through  $(\epsilon, G(\epsilon))$  with derivative  $G'(\epsilon)$  crosses the  $y$ -axis. It is clear that all four conditions of Theorem 8 are satisfied, and therefore  $\mathcal{F}$  is not testable with respect to  $\mathcal{X}$ .

EXAMPLE 20: SQUARE INTEGRABILITY OF A DENSITY IS NOT TESTABLE. Consider the class  $\mathcal{F}$  of all densities  $f$  for which  $\int f^2 dx < \infty$ . Then the same argument as in the previous example shows that  $\mathcal{F}$  is not testable with respect to  $\mathcal{X}$ , the class of monotonically decreasing densities on  $[0, \infty)$  that are continuous on  $(0, \infty)$  (and thus have possibly an infinite peak at the origin). For the construction involving continuations, just note that a density with  $\int f^2 = \infty$  is continued on  $[0, \epsilon)$  as in Example 19 by a bounded density (which necessarily has  $\int f^2 < \infty$ ). The other continuation argument is trickier and requires a density whose behavior near the origin is as  $c/2\sqrt{x}$  (and thus is not square integrable), and whose distribution function  $G$  evolves as  $c\sqrt{x}$ . By choice of  $c$ , such a continuation exists that meets the requirements of continuity and monotonicity as well.

EXAMPLE 21: CONTINUITY OF A DENSITY IS NOT TESTABLE. Let  $\mathcal{F}$  be the class of densities such that each  $f \in \mathcal{F}$  has a version which is continuous at 0. This class is not testable. To see this, we apply Theorem 8 in the following setup:  $\mathcal{X}$  is the class of all strictly monotonically increasing densities on  $[0, 1]$ , continuous on  $(0, \infty)$ , with  $f(1) = 1$  and  $f(x) = (2 - x)_+$ ,  $x > 1$ . Taking again  $S_\epsilon = (0, \epsilon)$ , it is easy to see that conditions (3) and (4) of Theorem 8 are satisfied, which implies the non-testability of  $\mathcal{F}$  with respect to  $\mathcal{X}$ . Indeed, we apply the same extension arguments from the previous examples, and note that the distribution functions  $G$  should be convex and increasing on  $[0, 1]$ , with continuous derivatives there, and with  $G'(0) = 0$  if and only if  $f$  is continuous at 0 (and  $f(0) = 0$ ). Thus, within the class, we may always continue a  $G$  with  $G'(0) > 0$  on  $[0, \epsilon]$  by one having  $G'(0) = 0$  using a quadratic Bezier spline having the specified derivatives at 0 and  $\epsilon$ . Similarly, a function  $G$  with  $G'(0) = 0$  may be continued on

$[0, \epsilon]$  by a quadratic Bezier spline  $B$  having  $B'(0) = G(\epsilon)/(2\epsilon) > 0$  and  $B'(\epsilon) = G'(\epsilon)$ . It should be easy to generalize and conclude that the class of densities with  $\int f^r < \infty$  (for fixed  $r > 1$ ) is not testable with respect to the decreasing densities on  $[0, 1]$ . The same goes for conditions like  $\int f \log^a(1 + f) < \infty$  for any  $a > 0$ , and indeed many other functionals.

EXAMPLE 22: LIPSCHITZ CONTINUITY OF A DENSITY IS NOT TESTABLE. Let  $\mathcal{F}$  be the class of densities such that each  $f \in \mathcal{F}$  has a version which is Lipschitz at 0. This class is not testable. To see this, we consider  $\mathcal{X}$  as in the previous example and ask additionally for continuity on the whole line (so that for each density,  $f(0) = 0$ ). We need to slightly modify the continuation argument. If  $G$  corresponds to a density in  $\mathcal{F} \cap \mathcal{X}$ , then, near the origin,  $G(x) \leq Cx^2$  for some constant  $C$ . Such a  $G$  may be continued on  $[0, \epsilon]$  by another  $G$  behaving as  $cx^{3/2}$  near the origin, where  $c > 0$  is picked appropriately, while maintaining the other restrictions imposed by  $\mathcal{X}$ . Note that the corresponding density behaves as  $\sqrt{x}$  near the origin, and is thus not Lipschitz. If  $G$  corresponds to a density in  $\mathcal{F}^c \cap \mathcal{X}$ , then, near the origin, we may continue  $G$  by another function within  $\mathcal{X}$  that behaves as  $Cx^2$  near the origin for an appropriately small but positive  $C$ . Thus, within  $\mathcal{X}$ , Lipschitz continuity at even one point is not testable.

We leave it to the reader as a simple exercise now to verify that we cannot test whether  $\mathbf{E}|X|^a < \infty$  for fixed  $a > 0$  with respect to the class of bounded symmetric unimodal densities (by continuations of the tails by finite support or heavy-tailed pieces). Similarly, the finiteness of a moment generating function cannot be tested.

## 11. Non-testability of mixture classes

Consider the class  $\mathcal{X}$  of densities which may be obtained as densities of random variables of the form  $X = F(U, Z)$ , where  $F : [0, 1] \times \{1, 2, \dots\} \rightarrow \mathcal{R}$  is a fixed measurable function,  $U$  is a uniform random variable on  $[0, 1]$ , and  $Z$  is an arbitrary positive-integer-valued random variable, independent of  $U$ . We assume that  $F$  is such that  $F(U, Z)$  has a density for all possible  $Z$ , and that for any two different distributions of  $Z$ ,  $F(U, Z)$  has different densities. Let  $\mathcal{F}$  be the subclass of  $\mathcal{X}$  containing all densities such that  $Z$  has a finite support.

THEOREM 9. *The class  $\mathcal{F}$  defined above is not testable with respect to  $\mathcal{X}$ .*

PROOF. We proceed similarly as in Theorem 8. Just like there, we need a coupling between samples drawn from different densities of  $\mathcal{X}$ . A simple way of doing it is as follows: Let  $U_1, V_1, U_2, V_2, \dots$  be i.i.d. uniform random variables on  $[0, 1]$ . Given a distribution  $(p_1, p_2, \dots)$  on the set of positive integers, let  $Z_i = j$  if and only if  $V_i \in \left[ \sum_{k=0}^{j-1} p_k, \sum_{k=0}^j p_k \right)$ . The sample drawn from a density  $f \in \mathcal{X}$  is now  $X_1, X_2, \dots$ , where  $X_i = F(U_i, Z_i)$ . The densities in  $\mathcal{F}$  are characterized by the distributions  $(p_1, p_2, \dots)$  which have finite support (i.e., only finitely many  $p_i$ 's are nonzero). Call this class  $\mathcal{P}$ . Similarly, the class  $\mathcal{X} - \mathcal{F}$  corresponds to the class  $\mathcal{Q}$  of distributions with infinite support.

Assume that the statement is false, and there exists a test  $T_n$  such that for any  $f \in \mathcal{F}$ , with probability one, there exists an integer  $N(\omega)$  such that

$$T_n(X_1, \dots, X_n) = 1 \quad \text{if } n > N(\omega),$$

and for any  $f \in \mathcal{X} - \mathcal{F}$ , with probability one, there exists an integer  $N(\omega)$  such that

$$T_n(X_1, \dots, X_n) = 0 \quad \text{if } n > N(\omega).$$

Let  $\delta_k = 2^{-k-2}$ ,  $k = 1, 2, \dots$

Let  $(p_1^{(1)}, p_2^{(1)}, \dots) \in \mathcal{P}$  be arbitrary. Then there exists an integer  $N_1 = N_1(\omega)$  such that

$$\mathbf{P} \left\{ T_n(X_1^{(1)}, \dots, X_n^{(1)}) = 1 \text{ for all } n \geq N_1 \right\} > 1 - \delta_1,$$

where  $X_1^{(1)}, \dots, X_n^{(1)}$  is the sample drawn from the corresponding density as described above. Next consider another distribution  $(p_1^{(2)}, p_2^{(2)}, \dots) \in \mathcal{Q}$  such that

$$\mathbf{P} \left\{ X_1^{(1)} = X_1^{(2)}, \dots, X_n^{(1)} = X_n^{(2)} \right\} > 1 - \delta_1.$$

Such a distribution may be constructed by choosing  $\epsilon_1 > 0$  such that  $(1 - \epsilon_1)^{N_1} > 1 - \delta_1$ , and defining  $p_i^{(2)} = p_i^{(1)}$  for all  $i < m$ , where  $m$  is the largest integer such that  $p_m^{(2)} > 0$ ,  $p_m^{(2)} = \min(0, p_m^{(1)} - \epsilon_1)$ , and  $p_i^{(2)} = 2^{-(i-m+1)} \min(\epsilon_1, p_m^{(1)})$  for all  $i \geq m$ .

This step may be iterated just like in Theorem 8, leading to the construction of a limiting density in  $\mathcal{X} - \mathcal{F}$  for which, with probability greater than  $1/2$ , the test  $T_n$  fails for every odd  $n$ . The proof is finished.  $\square$

EXAMPLE 23: NORMAL MIXTURES. It is not testable whether a normal mixture density has finitely or infinitely many components. This may be straightforwardly cast in the framework of Theorem 9: let  $\mathcal{X}$  be the class of all densities of the form

$$f(x) = \frac{1}{\sqrt{2\pi}} \sum_{i=1}^{\infty} \frac{p_i}{\sigma_i} e^{-\frac{(x-m_i)^2}{2\sigma_i^2}},$$

where  $p_1, p_2, \dots$  is a probability vector,  $\sigma_1, \sigma_2, \dots$  are positive numbers, and  $m_1, m_2, \dots$  are arbitrary real numbers. Assume that all parameters except the  $p_i$ 's are known, and that no two pairs  $(m_i, \sigma_i)$  are identical. This class is  $\mathcal{X}$ . According to Theorem 9, the class  $\mathcal{F}$  consisting of all densities of the above form such that  $p_i > 0$  for only finitely many  $i$ 's is not testable with respect to  $\mathcal{X}$ . From this, it follows certainly that the class of all finite mixtures of normal densities is not testable (here, the parameters  $m_i$  and  $\sigma_i$  are unrestricted).

EXAMPLE 24: CHARACTERISTIC FUNCTIONS WITH COMPACT SUPPORT. Let  $\mathcal{F}$  be the class of all densities whose characteristic function  $\varphi$  has bounded support. This class is not testable. To see this, we take densities  $f_n$  with characteristic function  $(1 - |t|/n)_+$ , and let  $\mathcal{X}$  be the class of all mixtures of finitely or

infinitely many components  $f_n$ . Then the class of finite mixtures of  $f_n$ 's is not testable with respect to  $\mathcal{X}$ , by Theorem 9.

EXAMPLE 25: CHARACTERISTIC FUNCTIONS WITH EXPONENTIALLY DECREASING TAILS. Let  $\mathcal{F}$  be the class of all densities whose characteristic function  $\varphi$  drops off exponentially quickly, i.e., for which  $|\varphi(t)| \leq Ce^{-a|t|}$  for positive constants  $C, a$ . This class is not testable. To see this, we take Cauchy densities  $f_n$  with characteristic function  $e^{-|t|/n}$ , and let  $\mathcal{X}$  be the class of all mixtures of finitely or infinitely many components  $f_n$ . Then the class of finite mixtures of  $f_n$ 's is not testable with respect to  $\mathcal{X}$ , by Theorem 9. However, for any infinite mixture, the mixture density does not have an exponentially decaying characteristic function.

## 12. References

- L. Birgé, “Non-asymptotic minimax risk for Hellinger balls,” *Probability and Mathematical Statistics*, vol. 5, pp. 21–29, 1985.
- L. Birgé, “On estimating a density using Hellinger distance and some other strange facts,” *Probability Theory and Related Fields*, vol. 71, pp. 271–291, 1986.
- T. M. Cover, “On determining the irrationality of the mean of a random variable,” *Annals of Statistics*, vol. 1, p. 862,, 1973.
- L. Devroye, “On arbitrarily slow rates of global convergence in density estimation,” *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, vol. 62, pp. 475–483, 1983.
- L. Devroye, “A simple algorithm for generating random variates with a log-concave density,” *Computing*, vol. 33, pp. 247–257, 1984.
- L. Devroye, “A note on the L1 consistency of variable kernel estimates,” *Annals of Statistics*, vol. 13, pp. 1041–1049, 1985.
- L. Devroye, *A Course In Density Estimation*, Birkhäuser Verlag, Boston, 1987.
- L. Devroye, “Asymptotic performance bounds for the kernel estimate,” *Annals of Statistics*, vol. 16, pp. 1162–1179, 1988.
- L. Devroye, “Nonparametric density estimates with improved performance on given sets of densities,” *Statistics (Mathematische Operationsforschung und Statistik)*, vol. 20, pp. 357–376, 1989.
- L. Devroye, “Universal smoothing factor selection in density estimation: theory and practice (with discussion),” *Test*, vol. 6, pp. 223–320, 1997.
- L. Devroye, L. Györfi, and G. Lugosi, *A Probabilistic Theory of Pattern Recognition*, Springer-Verlag, New York, 1996.
- L. Devroye and G. Lugosi, “A universally acceptable smoothing factor for kernel density estimation,” *Annals of Statistics*, vol. 24, pp. 2499–2512, 1996.

- L. Devroye and G. Lugosi, “Nonasymptotic universal smoothing factors, kernel complexity and Yatracos classes,” *Annals of Statistics*, vol. 25, pp. 2626–2637, 1997.
- L. Devroye, G. Lugosi, and F. Udina, “Inequalities for a new data-based method for selecting nonparametric densities,” Technical Report, Facultat de Ciències Econòmiques, Universitat Pompeu Fabra, 1998.
- W. Hoeffding, “Probability inequalities for sums of bounded random variables,” *Journal of the American Statistical Association*, vol. 58, pp. 13–30, 1963.
- J. Kopolowitz, “Abstracts of papers,” in: *International Symposium on Information Theory*, p. 64, Cornell University, Ithaca, NY, 1977.
- S. R. Kulkarni and O. Zeitouni, “A general classification rule for probability measures,” *Annals of Statistics*, vol. 23, pp. 1393–1407, 1995.
- G. G. Lorentz, *Approximation of Functions*, Holt, Rinehart and Winston, New York, 1966.
- P. Massart, “The tight constant in the Dvoretzky-Kiefer-Wolfowitz inequality,” *Annals of Probability*, vol. 18, pp. 1269–1283, 1990.
- C. McDiarmid, “On the method of bounded differences,” in: *Surveys in Combinatorics*, ed. J. Siemons, vol. 141, pp. 148–188, London Mathematical Society Lecture Note Series, Cambridge University Press, 1989.
- H. L. Royden, *Real Analysis*, Macmillan Publishing Co., New York, 1968.
- V. N. Vapnik and A. Ya. Chervonenkis, “On the uniform convergence of relative frequencies of events to their probabilities,” *Theory of Probability and its Applications*, vol. 16, pp. 264–280, 1971.
- R. L. Wheeden and A. Zygmund, *Measure and Integral*, Marcel Dekker, New York, 1977.
- Y. G. Yatracos, “Rates of convergence of minimum distance estimators and Kolmogorov’s entropy,” *Annals of Statistics*, vol. 13, pp. 768–774, 1985.
- Y. G. Yatracos, “A note on  $L_1$  consistent estimation,” *Canadian Journal of Statistics*, vol. 16, pp. 283–292, 1988.